



STIC Search Report

Biotech-Chem Library

STIC Database Tracking Number: 113210

TO: Emily M Le
Location: Rem 3c75
Tuesday, February 03, 2004 3041
Art Unit: 1648
Phone: 272-0903
Serial Number: 10 / 600361

From: Jan Delaval
Location: Biotech-Chem Library
Remsen Building – 1A51
Phone: 272-2504

jan.delaval@uspto.gov

Search Notes

113210

Delaval, Jan

From: Le, Emily
Sent: Friday, January 30, 2004 3:36 PM
To: Delaval, Jan
Subject: RE: text search: 10/600361

Lo siento! 10/600361

-----Original Message-----

From: Delaval, Jan
Sent: Friday, January 30, 2004 3:34 PM
To: Le, Emily
Subject: RE: text search

Emily -

Serial number, please! Merci!

-----Original Message-----

From: Le, Emily
Sent: Friday, January 30, 2004 3:33 PM
To: Delaval, Jan
Subject: text search

Jan,

please provide a text search of the following: dendritic cells AND inactivated human immunodeficiency virus (HIV). Thanks!

Emily Le
Remsen, 3C35
(571) 272-0903

SEARCH REQUEST FORM

11/3/210

Requestor's Name: _____ Serial Number: _____
Date: _____ Phone: _____ Art Unit: _____

Search Topic:

Please write a detailed statement of search topic. Describe specifically as possible the subject matter to be searched. Define any terms that may have a special meaning. Give examples or relevant citations, authors, keywords, etc., if known. For sequences, please attach a copy of the sequence. You may include a copy of the broadest and/or most relevant claim(s).

STAFF USE ONLY

Date completed: 2/3/21
Searcher: _____
Terminal time: _____
Elapsed time: 1:15
CPU time: _____
Total time: _____
Number of Searches: _____
Number of Databases: _____

Search Site
____ STIC
____ CM-1
____ Pre-S
Type of Search
____ N.A. Sequence
____ A.A. Sequence
____ Structure
____ ☒ Bibliographic

Vendors
____ IG
____ ☒ STN
____ Dialog
____ APS
____ Geninfo
____ SDC
____ DARC/Questel
____ Other

BEST AVAILABLE COPY

=> d his

(FILE 'HOME' ENTERED AT 10:36:24 ON 03 FEB 2004)
SET COST OFF

FILE 'HCAPLUS' ENTERED AT 10:36:34 ON 03 FEB 2004

E DENDRITIC CELL/CT
E E3+ALL
L1 529 S E8,E9
L2 7326 S E7
L3 45 S E13
L4 7836 S L1-L3
L5 11672 S DENDRITIC CELL
L6 11672 S L4,L5
E HIV/CT
E E3+ALL
L7 9029 S E2
E E6+ALL
L8 12470 S E7,E8,E9,E10
L9 19620 S E6
E E5+ALL
L10 16484 S E6
L11 36735 S E5+NT
L12 635 S L6 AND L7-L11
L13 639 S L6 AND HIV
L14 613 S L6 AND HUMAN IMMUNODEFICIEN? VIRUS
L15 798 S L12-L14
L16 32 S L15 AND INACTIV?
L17 58 S L15 AND PULS?
L18 7 S L16 AND L17
E CD8/CT
E E10+ALL
L19 8218 S E20
L20 84 S L15 AND L19
L21 150 S L15 AND CD8
L22 150 S L20,L21
L23 20 S L22 AND L16,L17
L24 3 S L18 AND L23
L25 4 S L18 NOT L24
L26 7 S L24,L25
L27 17 S L23 NOT L26
SEL DN AN 6 14 15 16 17
L28 5 S E1-E15 AND L27
L29 12 S L26,L28
L30 35 S L17 NOT L23-L29
SEL DN AN 25
L31 1 S E16-E18
L32 13 S L29,L31
L33 1 S US20040009194/PN OR US2002-390625#/AP, PRN
E ANDRIEU J/AU
L34 95 S E3,E6,E7,E12,E13,E17
E LU L/AU
L35 437 S E3-E26
E LU LOUIS/AU
L36 5 S E3,E4
L37 4 S L15 AND L34-L36
L38 4 S L34 AND L35-L36
L39 19 S L32,L33,L37,L38
L40 15 S L39 AND ?ACTIV?
L41 4 S L39 NOT L40

=> fil hcaplus

FILE 'HCAPLUS' ENTERED AT 10:59:40 ON 03 FEB 2004

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2004 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 3 Feb 2004 VOL 140 ISS 6
FILE LAST UPDATED: 2 Feb 2004 (20040202/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> d 140 all tot

L40 ANSWER 1 OF 15 HCAPLUS COPYRIGHT 2004 ACS on STN
AN 2004:39604 HCAPLUS
ED Entered STN: 16 Jan 2004
TI Methods, and compositions for a therapeutic antigen presenting cell vaccine for treatment of immunodeficiency virus
IN **Andrieu, Jean-Marie; Lu, Louis**
PA Fr.
SO U.S. Pat. Appl. Publ., 29 pp.
CODEN: USXXCO
DT Patent
LA English
IC ICM A61K039-21
ICS A61K031-551
NCL 424208100; 514220000
CC 63 (Pharmaceuticals)
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 2004009194	A1	20040115	US 2003-600361	20030620 <--
PRAI	US 2002-390625P	P	20020621	<--	

AB One aspect of this invention provides a composition capable of eliciting an immune response to an immunodeficiency virus in mammals, wherein the composition is comprised of an **inactivated** virus-pulsed antigen presenting cell. In another aspect the aforementioned composition may also contain a combination of an **inactivated** virus-pulsed antigen presenting cell and an immunodeficiency protease inhibitor. Still other aspects of this invention provide for methods of treating mammals with an **inactivated** virus-pulsed antigen presenting cell, the vaccines related to such cells.

L40 ANSWER 2 OF 15 HCAPLUS COPYRIGHT 2004 ACS on STN
AN 2003:747940 HCAPLUS
DN 139:394729
ED Entered STN: 24 Sep 2003
TI Presentation of Exogenous Whole **Inactivated** Simian Immunodeficiency Virus by Mature **Dendritic Cells** Induces CD4+ and CD8+ T-cell Responses
AU Frank, Ines; Santos, John J.; Mehlhop, Erin; Villamide-Herrera, Loreley; Santisteban, Christine; Gettie, Agegnehu; Ignatius, Ralf; Lifson, Jeffrey

- D.; Pope, Melissa
 CS Population Council, Center for Biomedical Research, New York, NY, USA
 SO JAIDS, Journal of Acquired Immune Deficiency Syndromes (2003), 34(1), 7-19
 CODEN: JJASFJ; ISSN: 1525-4135
 PB Lippincott Williams & Wilkins
 DT Journal
 LA English
 CC 15-8 (Immunochemistry)
 AB Interactions between **HIV-1** and **dendritic cells**
 (DCs) play an important role in the initial establishment and spread of infection and development of antiviral immunity. The authors used chemical **inactivated** aldrithiol-2 (AT-2) simian immunodeficiency virus (SIV) with functional envelope glycoproteins to study virus interactions with DCs and developed an in vitro system to evaluate the quality of SIV antigen (Ag) presentation by DCs to T cells. AT-2 SIV interacts authentically with T cells and DCs and thus allows assessment of natural SIV-specific responses. CD4+ and **CD8+** T cells from blood or lymph nodes of SIV-infected macaques released interferon- γ (IFN γ) and proliferated in response to a variety of AT-2 SIV isolates. Responses did not vary significantly as a function of the quant. envelope glycoprotein content of the virions. Presentation of Ags derived from AT-2 SIV by DCs was more potent than presentation by comparably Ag-loaded monocytes. Interestingly, **SIV-pulsed** mature DCs stimulated both CD4+ and **CD8+** T-cell responses, whereas immature DCs primarily stimulated CD4+ T cells. Further studies using AT-2 **inactivated** virus may help to define better the details of the virus-DC interactions critical for infection vs. induction of antiviral immune responses.
 ST simian immunodeficiency virus **dendritic cell** CD4
 CD8 T lymphocyte; antigen presentation interferon
 IT Cell proliferation
 (T cell; presentation of exogenous whole **inactivated** simian immunodeficiency virus by mature **dendritic cells** induces CD4+ and **CD8+** T-cell responses)
 IT Antigen presentation
 CD4-positive T cell
 CD8-positive T cell
 Dendritic cell
 Macaca mulatta
 Monocyte
 Simian immunodeficiency virus
 (presentation of exogenous whole **inactivated** simian immunodeficiency virus by mature **dendritic cells** induces CD4+ and **CD8+** T-cell responses)
 IT T cell (lymphocyte)
 (proliferation; presentation of exogenous whole **inactivated** simian immunodeficiency virus by mature **dendritic cells** induces CD4+ and **CD8+** T-cell responses)
 IT Interferons
 RL: BSU (Biological study, unclassified); BIOL (Biological study)
 (γ ; presentation of exogenous whole **inactivated** simian immunodeficiency virus by mature **dendritic cells** induces CD4+ and **CD8+** T-cell responses)
 RE.CNT 59 THERE ARE 59 CITED REFERENCES AVAILABLE FOR THIS RECORD
 RE
 (1) Allaway, G; AIDS Res Hum Retroviruses 1995, V11, P533 HCAPLUS
 (2) Arthur, L; AIDS Res Hum Retroviruses 1998, V14(Suppl 3), PS311
 (3) Banchereau, J; Nature 1998, V392, P245 HCAPLUS
 (4) Barouch, D; J Immunol 2002, V168, P562 HCAPLUS
 (5) Barratt-Boyes, S; J Immunol 2000, V164, P2487 HCAPLUS
 (6) Benveniste, R; J Med Primatol 1990, V19, P351 HCAPLUS
 (7) Boes, M; Nature 2002, V418, P983 HCAPLUS
 (8) Buseyne, F; Nat Med 2001, V7, P344 HCAPLUS

- (9) Chertova, E; Curr Mol Med 2003, V3, P265 HCAPLUS
 - (10) Chertova, E; J Virol 2002, V76, P5315 HCAPLUS
 - (11) Chow, A; Nature 2002, V418, P988 HCAPLUS
 - (12) Connor, R; J Virol 1998, V72, P7501 HCAPLUS
 - (13) Desrosiers, R; AIDS 1995, V9(Suppl A), PS137
 - (14) Douek, D; Nature 2002, V417, P95 HCAPLUS
 - (15) Engelmayer, J; J Immunol 1999, V163, P6762 HCAPLUS
 - (16) Esser, M; J Virol 2001, V75, P1152 HCAPLUS
 - (17) Esser, M; J Virol 2001, V75, P6173 HCAPLUS
 - (18) Fayette, J; J Exp Med 1997, V185, P1909 HCAPLUS
 - (19) Feuerstein, B; J Immunol Methods 2000, V245, P15 HCAPLUS
 - (20) Frank, I; Curr Mol Med 2002, V2, P229 HCAPLUS
 - (21) Frank, I; Immunobiology 2001, V204, P622 HCAPLUS
 - (22) Frank, I; J Virol 2002, V76, P2936 HCAPLUS
 - (23) Garrett, W; Cell 2000, V102, P325 HCAPLUS
 - (24) Gaschen, B; Science 2002, V296, P2354 HCAPLUS
 - (25) Granelli-Piperno, A; J Immunol 2000, V165, P6620 HCAPLUS
 - (26) Ho, D; Cell 2002, V110, P135 HCAPLUS
 - (27) Horton, H; J Virol 2002, V76, P7187 HCAPLUS
 - (28) Hu, J; AIDS Res Hum Retroviruses 1999, V15, P1305 MEDLINE
 - (29) Hu, J; Lab Invest 1998, V78, P435 MEDLINE
 - (30) Ignatius, R; J Med Primatol 1998, V27, P121 HCAPLUS
 - (31) Ignatius, R; J Virol 2000, V74, P11329 HCAPLUS
 - (32) Labranche, C; J Virol 1995, V69, P5217 HCAPLUS
 - (33) Larsson, M; AIDS 2002, V16, P1319 HCAPLUS
 - (34) Larsson, M; AIDS 2002, V16, P171
 - (35) Letvin, N; Annu Rev Immunol 2002, V20, P73 HCAPLUS
 - (36) Lifson, J; J Med Primatol 2002, V31, P205 HCAPLUS
 - (37) Litinskiy, M; Nat Immunol 2002, V3, P822 HCAPLUS
 - (38) Lu, W; Nat Med 2002, V9, P27
 - (39) Marovich, M; J Infect Dis 2002, V186, P1242 HCAPLUS
 - (40) Mascola, J; Vaccine 2002, V20, P1922 HCAPLUS
 - (41) McDonald, D; Science 2003, V300, P1295 HCAPLUS
 - (42) McMichael, A; Nat Rev Immunol 2002, V2, P283 HCAPLUS
 - (43) Mehlhop, E; J Immunol Methods 2002, V260, P219 HCAPLUS
 - (44) Nabel, G; Vaccine 2002, V20, P1945 HCAPLUS
 - (45) O'Doherty, U; J Immunol Methods 1997, V207, P185 HCAPLUS
 - (46) Peters, B; Vaccine 2002, V20, P688
 - (47) Pomerantz, R; Curr Opin Invest Drugs 2002, V3, P1133 HCAPLUS
 - (48) Pope, M; AIDS Res Hum Retroviruses 1997, V13, P819 MEDLINE
 - (49) Pope, M; Curr Mol Med 2003, V3, P229 HCAPLUS
 - (50) Randolph, G; J Exp Med 2002, V196, P517 HCAPLUS
 - (51) Randolph, G; Semin Immunol 2001, V13, P267 HCAPLUS
 - (52) Rossio, J; J Virol 1998, V72, P7992 HCAPLUS
 - (53) Santini, S; J Exp Med 2000, V191, P1777 HCAPLUS
 - (54) Steinman, R; J Clin Invest 2002, V109, P1519 HCAPLUS
 - (55) Teleshova, N; J Leuk Biol in press 2003
 - (56) Trombetta, E; Science 2003, V299, P1400 HCAPLUS
 - (57) Watts, C; Annu Rev Immunol 1997, V15, P821 HCAPLUS
 - (58) Willey, R; J Virol 2003, V77, P1163 HCAPLUS
 - (59) Zhu, T; J Antimicrob Chemother 2002, V50, P309 HCAPLUS
- L40 ANSWER 3 OF 15 HCAPLUS COPYRIGHT 2004 ACS on STN
- AN 2003:620148 HCAPLUS
- DN 139:196131
- ED Entered STN: 13 Aug 2003
- TI Induction of protective immune responses against R5 human immunodeficiency virus type 1 (HIV-1) infection in hu-PBL-SCID mice by intrasplenic immunization with HIV-1-pulsed dendritic cells:
- Possible involvement of a novel factor of human CD4+ T-cell origin
- AU Yoshida, Atsushi; Tanaka, Reiko; Murakami, Tsutomu; Takahashi, Yoshiaki; Koyanagi, Yoshio; Nakamura, Masataka; Ito, Mamoru; Yamamoto, Naoki;

Tanaka, Yuetsu

CS Department of Immunology, Graduate School and Faculty of Medicine,
University of the Ryukyus, Okinawa, 903-0215, Japan

SO Journal of Virology (2003) 77(16), 8719-8728

CODEN: JOVIAM; ISSN: 0022-538X

PB American Society for Microbiology

DT Journal

LA English

CC 15-8 (Immunochemistry)

Section cross-reference(s): 1, 63

AB The potential of a **dendritic cell** (DC)-based vaccine against **human immunodeficiency virus** type 1 (HIV-1) infection in humans was explored with SCID mice reconstituted with human peripheral blood mononuclear cells (PBMC). HIV-1-neg. normal human PBMC were transplanted directly into the spleens of SCID mice (hu-PBL-SCID-spl mice) together with autologous mature DCs **pulsed** with either **inactivated HIV** -1 (strain R5 or X4) or ovalbumin (OVA), followed by a booster injection 5 days later with autologous DCs **pulsed** with the same resp. antigens. Five days later, these mice were challenged i.p. with R5 HIV-1JR-CSF. Anal. of infection at 7 days postinfection showed that the DC-HIV-1-immunized hu-PBL-SCID-spl mice, irres. of the HIV-1 isolate used for immunization, were protected against HIV-1 infection. In contrast, none of the DC-OVA-immunized mice were protected. Sera from the DC-HIV-1- but not the DC-OVA-immunized mice inhibited the in vitro infection of **activated** PBMC and macrophages with R5, but not X4, HIV -1. Upon restimulation with HIV-1 in vitro, the human CD4+ T cells derived from the DC-HIV-1-immunized mice produced a similar R5 HIV-1 suppressor factor. Neutralizing antibodies against human RANTES, MIP-1 α , MIP-1 β , alpha interferon (IFN- α), IFN- β , IFN- γ , interleukin-4 (IL-4), IL-10, IL-13, IL-16, MCP-1, MCP-3, tumor necrosis factor alpha (TNF- α), or TNF- β failed to reverse the HIV-1-suppressive **activity**. These results show that **inactivated HIV-1-pulsed** autologous DCs can stimulate splenic resident human CD4+ T cells in hu-PBL-SCID-spl mice to produce a yet-to-be-defined, novel soluble factor(s) with protective properties against R5 HIV-1 infection.

ST AIDS vaccine spleen **dendritic cell** CD4 T lymphocyte factor

IT Vaccines

(AIDS; a new factor involved in the induction of anti-HIV responses in hu-PBL-SCID mice by intrasplenic immunization with **HIV-1-pulsed dendritic cells**)

IT Chemokine receptors

RL: ADV (Adverse effect, including toxicity); BSU (Biological study, unclassified); BIOL (Biological study)

(CCR5; a new factor involved in the induction of anti-HIV responses in hu-PBL-SCID mice by intrasplenic immunization with **HIV-1-pulsed dendritic cells**)

IT Cell activation

(T cell; a new factor involved in the induction of anti-HIV responses in hu-PBL-SCID mice by intrasplenic immunization with **HIV-1-pulsed dendritic cells**)

IT CD4-positive T cell

Dendritic cell

Human

Human immunodeficiency virus 1

Macrophage

Mononuclear cell (leukocyte)

Spleen

(a new factor involved in the induction of anti-HIV responses

- in hu-PBL-SCID mice by intrasplenic immunization with **HIV-1-pulsed dendritic cells**)
- IT T cell (lymphocyte)
(**activation**; a new factor involved in the induction of anti-HIV responses in hu-PBL-SCID mice by intrasplenic immunization with **HIV-1-pulsed dendritic cells**)
- IT Proteins
RL: BSU (Biological study, unclassified); BIOL (Biological study) (from **HIV-1** infected CD4+ T lymphocytes with anti-HIV1 **activity**; a new factor involved in the induction of anti-HIV responses in hu-PBL-SCID mice by intrasplenic immunization with **HIV-1-pulsed dendritic cells**)
- IT Anti-AIDS agents
(vaccines; a new factor involved in the induction of anti-HIV responses in hu-PBL-SCID mice by intrasplenic immunization with **HIV-1-pulsed dendritic cells**)

RE.CNT 37 THERE ARE 37 CITED REFERENCES AVAILABLE FOR THIS RECORD
RE

- (1) Aaberge, I; Clin Exp Immunol 1996, V105, P12 HCAPLUS
- (2) Adachi, A; J Virol 1986, V59, P284 MEDLINE
- (3) Bombil, F; Immunobiology 1996, V195, P360 HCAPLUS
- (4) Delhem, N; J Immunol 1998, V161, P2060 HCAPLUS
- (5) Dhawan, S; J Leukoc Biol 1995, V58, P713 HCAPLUS
- (6) Geiben-Lynn, R; J Virol 2001, V75, P8306 HCAPLUS
- (7) Ifversen, P; Immunology 1995, V84, P111 HCAPLUS
- (8) Ito, M; Blood 2002, V100, P3175 HCAPLUS
- (9) Koyanagi, Y; J Virol 1997, V71, P2417 HCAPLUS
- (10) Koyanagi, Y; Science 1987, V236, P819 MEDLINE
- (11) Lane, B; J Immunol 1999, V163, P3653 HCAPLUS
- (12) Lapenta, C; Virology 1999, V263, P78 HCAPLUS
- (13) Lehner, T; J Immunol 1994, V153, P1858 HCAPLUS
- (14) Lehner, T; Nat Med 1996, V2, P767 HCAPLUS
- (15) Levy, J; Immunol Today 1996, V17, P217 HCAPLUS
- (16) Lu, W; Nat Med 2003, V9, P27 HCAPLUS
- (17) Mazingue, C; Eur J Immunol 1991, V21, P1763 MEDLINE
- (18) McCune, J; Annu Rev Immunol 1991, V9, P399 HCAPLUS
- (19) McNeely, T; Blood 1997, V90, P1141 HCAPLUS
- (20) Mengozzi, M; Proc Natl Acad Sci USA 2001, V98, P11644 HCAPLUS
- (21) Montaner, L; Science 1995, V267, P538 HCAPLUS
- (22) Mosier, D; Adv Immunol 1996, V63, P79 HCAPLUS
- (23) Mosier, D; Science 1991, V251, P791 MEDLINE
- (24) Ohteki, T; J Exp Med 1999, V189, P1981 HCAPLUS
- (25) Riley, J; J Immunol 1997, V158, P5545 HCAPLUS
- (26) Rizza, P; J Virol 1996, V70, P7958 HCAPLUS
- (27) Rossio, J; J Virol 1998, V72, P7992 HCAPLUS
- (28) Santini, S; J Exp Med 2000, V191, P1777 HCAPLUS
- (29) Saphire, A; J Virol 2001, V75, P9187 HCAPLUS
- (30) Shioda, T; Nature 1991, V349, P167 HCAPLUS
- (31) Subramani, S; Mol Cell Biol 1981, V1, P854 HCAPLUS
- (32) Suzuki, Y; J Virol 1999, V73, P316 HCAPLUS
- (33) Tanaka, T; J Exp Med 1993, V178, P1103 HCAPLUS
- (34) Tary-Lehmann, M; Immunol Today 1995, V16, P529 HCAPLUS
- (35) Zaitseva, M; Blood 2000, V96, P3109 HCAPLUS
- (36) Zhang, L; Science 2002, V26, P26
- (37) Zhao, X; J Virol 2002, V76, P3007 HCAPLUS

L40 ANSWER 4 OF 15 HCAPLUS COPYRIGHT 2004 ACS on STN
AN 2003:588083 HCAPLUS
DN 139:212744
ED Entered STN: 31 Jul 2003
TI Potent immune response against **HIV-1** and protection from virus

challenge in hu-PBL-SCID mice immunized with **inactivated virus-pulsed dendritic cells** generated in the presence of IFN- α

AU Lapenta, Caterina; Santini, Stefano M.; Logozzi, Mariantonia; Spada, Massimo; Andreotti, Mauro; Di Pucchio, Tiziana; Parlato, Stefania; Belardelli, Filippo
CS Laboratory of Virology, Istituto Superiore di Sanita, Rome, 00161, Italy
SO Journal of Experimental Medicine (2003), 198(2), 361-367
CODEN: JEMEAV; ISSN: 0022-1007
PB Rockefeller University Press
DT Journal
LA English
CC 15-8 (Immunochemistry)

AB A major challenge of AIDS research is the development of therapeutic vaccine strategies capable of inducing the humoral and cellular arms of the immune responses against **HIV-1**. In this work, the authors evaluated the capability of DCs **pulsed** with aldrithiol-2-**inactivated HIV-1** in inducing a protective antiviral human immune response in SCID mice reconstituted with human PBL (hu-PBL-SCID mice). Immunization of hu-PBL-SCID mice with DCs generated after exposure of monocytes to GM-CSF/IFN- α (IFN-DCs) and **pulsed** with **inactivated HIV-1** resulted in a marked induction of human anti-**HIV-1** antibodies, which was associated with the detection of anti-**HIV** neutralizing **activity** in the serum. This vaccination schedule also promoted the generation of a human **CD8+** T cell response against **HIV-1**, as measured by IFN- γ Elispot anal. Notably, when the hu-PBL-SCID mice immunized with antigen-**pulsed** IFN-DCs were infected with **HIV-1**, inhibition of virus infection was observed as compared with control animals. These results suggest that IFN-DCs **pulsed** with **inactivated HIV-1** can represent a valuable approach of immune intervention in **HIV-1**-infected patients.

ST HIV1 vaccine dendritic cell interferon

IT Vaccines

(AIDS; immune response against **HIV-1** and protection from virus challenge in hu-PBL-SCID mice immunized with **inactivated virus-pulsed dendritic cells** generated in presence of IFN- α)

IT **CD8-positive T cell**

Dendritic cell

Human

Human immunodeficiency virus 1

Lymph node

Spleen

(immune response against **HIV-1** and protection from virus challenge in hu-PBL-SCID mice immunized with **inactivated virus-pulsed dendritic cells** generated in presence of IFN- α)

IT Antibodies

RL: BSU (Biological study, unclassified); BIOL (Biological study) (neutralizing; immune response against **HIV-1** and protection from virus challenge in hu-PBL-SCID mice immunized with **inactivated virus-pulsed dendritic cells** generated in presence of IFN- α)

IT Anti-AIDS agents

(vaccines; immune response against **HIV-1** and protection from virus challenge in hu-PBL-SCID mice immunized with **inactivated virus-pulsed dendritic cells** generated in presence of IFN- α)

IT Interferons

RL: BSU (Biological study, unclassified); BIOL (Biological study) (α ; immune response against **HIV-1** and protection from

virus challenge in hu-PBL-SCID mice immunized with **inactivated virus-pulsed dendritic cells** generated in presence of IFN- α)

IT Interferons

RL: BSU (Biological study, unclassified); BIOL (Biological study) (γ ; immune response against **HIV-1** and protection from virus challenge in hu-PBL-SCID mice immunized with **inactivated virus-pulsed dendritic cells** generated in presence of IFN- α)

IT 83869-56-1, Granulocyte-macrophage colony-stimulating factor

RL: BSU (Biological study, unclassified); BIOL (Biological study) (immune response against **HIV-1** and protection from virus challenge in hu-PBL-SCID mice immunized with **inactivated virus-pulsed dendritic cells** generated in presence of IFN- α)

RE.CNT 27 THERE ARE 27 CITED REFERENCES AVAILABLE FOR THIS RECORD
RE

- (1) Banchereau, J; Annu Rev Immunol 2000, V18, P767 HCAPLUS
- (2) Borrow, P; J Virol 1994, V68, P6103 HCAPLUS
- (3) Brander, C; J Clin Invest 1998, V101, P2559 HCAPLUS
- (4) Brander, C; J Virol 1999, V73, P10191
- (5) Buseyne, F; Nat Med 2001, V7, P344 HCAPLUS
- (6) Chun, T; Nature 1999, V401, P874 HCAPLUS
- (7) de Jong, M; AIDS 1997, V11, PF79 MEDLINE
- (8) Finzi, D; Nat Med 1999, V5, P512 HCAPLUS
- (9) Gauduin, M; Nat Med 1997, V3, P1389 MEDLINE
- (10) Hart, D; Pathology 2001, V33, P479 MEDLINE
- (11) Jin, X; J Exp Med 1999, V189, P991 HCAPLUS
- (12) Jubault, V; AIDS 1998, V12, P2358 HCAPLUS
- (13) Le Bon, A; Immunity 2001, V14, P461 HCAPLUS
- (14) Lu, W; J Virol 2001, V75, P8949 HCAPLUS
- (15) Lu, W; Nat Med 2003, V9, P27 HCAPLUS
- (16) Muster, T; J Virol 1994, V68, P4031 HCAPLUS
- (17) Nehete, P; Vaccine 2001, V20, P813 HCAPLUS
- (18) Ogg, G; Science 1998, V279, P2103 HCAPLUS
- (19) Okamoto, Y; J Immunol 1998, V160, P69 HCAPLUS
- (20) Pantaleo, G; Annu Rev Microbiol 1996, V50, P825 HCAPLUS
- (21) Poignard, P; Immunity 1999, V10, P431 HCAPLUS
- (22) Proietti, E; J Immunol 2002, V169, P375 HCAPLUS
- (23) Rizza, P; J Virol 1996, V70, P7958 HCAPLUS
- (24) Santini, S; J Exp Med 2000, V191, P1777 HCAPLUS
- (25) Santodonato, L; J Immunol 2003, V170, P5195 HCAPLUS
- (26) Steinman, R; Int J Cancer 2001, V94, P459 HCAPLUS
- (27) Wagner, G; AIDS Res Hum Retroviruses 2002, V18, P143

L40 ANSWER 5 OF 15 HCAPLUS COPYRIGHT 2004 ACS on STN

AN 2003:221333 HCAPLUS

DN 138:236437

ED Entered STN: 21 Mar 2003

TI **Dendritic cells** and the promise of therapeutic vaccines for **human immunodeficiency virus (HIV)-1**

AU Walsh, Stephen R.; Bhardwaj, Nina; Gandhi, Rajesh T.

CS Division of Infectious Diseases, Department of Medicine, Massachusetts General Hospital, Boston, MA, 02114, USA

SO Current HIV Research (2003), 1(2), 205-216

CODEN: CHRUBF; ISSN: 1570-162X

PB Bentham Science Publishers Ltd.

DT Journal; General Review

LA English

CC 15-0 (Immunochemistry)

AB A review. Treatment of **human immunodeficiency virus (HIV)-1** infection with potent antiretroviral

medications has provided considerable clin. benefit. However because of the limitations of current therapy, innovative approaches are needed to better control HIV-1 infection. Several studies have suggested that robust CD4+ T helper and CD8+ T cell responses may contribute to the immunol. control of HIV-1 infection in certain individuals. Most chronically infected patients, however, cannot control the infection and may benefit from stimulation of cellular immunity with immunotherapy. **Dendritic cells** (DCs) are potent professional antigen-presenting cells (APCs) and have a central role in directing the adaptive immune response to pathogens. The ability of DCs to stimulate naive T cells has long been thought to be crucial in initiating an effective immune response. As DCs are uniquely situated at the interface between the innate and adaptive immune systems, they are currently under intense scrutiny as potential adjuvants for vaccines in many clin. settings. Studies in healthy volunteers and patients with cancer have shown that antigen-pulsed DCs can boost both CD8+ and CD4+ T cell responses in vivo. Based on these promising findings, ex vivo antigen-pulsed DCs are being **actively** investigated in studies aimed at developing a therapeutic vaccine for individuals with HIV-1 infection.

ST review **dendritic cell** vaccine HIV1 cytokine antigen

IT Vaccines

(AIDS; **dendritic cells** and the promise of therapeutic vaccines for HIV-1)

IT Immunostimulants

(adjuvants; **dendritic cells** and the promise of therapeutic vaccines for HIV-1)

IT CD4-positive T cell

CD8-positive T cell

Dendritic cell

Human

Human immunodeficiency virus 1

(**dendritic cells** and the promise of therapeutic vaccines for HIV-1)

IT Cytokines

RL: BSU (Biological study, unclassified); BIOL.(Biological study)

(**dendritic cells** and the promise of therapeutic vaccines for HIV-1)

IT Antigens

RL: BSU (Biological study, unclassified); PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(**dendritic cells** and the promise of therapeutic vaccines for HIV-1)

IT Vaccines

(tumor; **dendritic cells** and the promise of therapeutic vaccines for HIV-1)

IT Anti-AIDS agents

Antitumor agents

(vaccines; **dendritic cells** and the promise of therapeutic vaccines for HIV-1)

RE.CNT 159 THERE ARE 159 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

- (1) Albert, M; Nature 1998, V392, P86 HCAPLUS
- (2) Allen, T; Trends in Immunology 2002, V23, P456 HCAPLUS
- (3) Altfeld, M; Journal of Experimental Medicine 2001, V193, P169 HCAPLUS
- (4) Andrieu, M; 9th Conference on Retroviruses and Opportunistic Infections 2002
- (5) Autran, B; Science 1997, V277, P112 HCAPLUS
- (6) Banchereau, J; Annual Review of Immunology 2000, V18, P767 HCAPLUS
- (7) Banchereau, J; Cancer Research 2001, V61, P6451 HCAPLUS
- (8) Banchereau, J; Nature 1998, V392, P245 HCAPLUS
- (9) Baribaud, F; Virology 2001, V286, P1 HCAPLUS
- (10) Bell, D; Journal of Experimental Medicine 1999, V190, P1417 HCAPLUS

- (11) Belz, G; Journal of Experimental Medicine 2002, V196, P1099 HCAPLUS
- (12) Bender, A; Journal of Immunological Methods 1996, V196, P121 HCAPLUS
- (13) Bhardwaj, N; Nature Medicine 2003, V9, P13 HCAPLUS
- (14) Bhardwaj, N; Trends in Molecular Medicine 2001, V7, P388 HCAPLUS
- (15) Blankson, J; Annual Review of Medicine 2002, V53, P557 HCAPLUS
- (16) Burch, P; Clinical Cancer Research 2000, V6, P2175 HCAPLUS
- (17) Cameron, P; Science 1992, V257, P383 MEDLINE
- (18) Carcelain, G; Journal of Virology 2001, V75, P234 HCAPLUS
- (19) Carr, A; Lancet 2000, V356, P1423 HCAPLUS
- (20) Cella, M; Current Opinion in Immunology 1997, V9, P10 HCAPLUS
- (21) Cella, M; Journal of Experimental Medicine 1996, V184, P747 HCAPLUS
- (22) Chan, R; Immunology Letters 2002, V83, P101 HCAPLUS
- (23) Chen, Q; International Journal of Cancer 1994, V56, P755 HCAPLUS
- (24) Dallal, R; Current Opinion in Immunology 2000, V12, P583 HCAPLUS
- (25) Davey, R; Proceedings of the National Academy of Sciences 1999, V96, P15109 HCAPLUS
- (26) Davis, N; Journal of Virology 2000, V74, P371 HCAPLUS
- (27) Dhodapkar, M; Journal of Clinical Immunology 2000, V20, P167 HCAPLUS
- (28) Dhodapkar, M; Journal of Clinical Investigation 1999, V104, P173 HCAPLUS
- (29) Dhodapkar, M; Journal of Clinical Investigation 2000, V105, PR9 HCAPLUS
- (30) Dhodapkar, M; Journal of Experimental Medicine 2001, V193, P233 HCAPLUS
- (31) Eggert, A; Cancer Research 1999, V59, P3340 HCAPLUS
- (32) Engelmayer, J; Journal of Virology 2001, V75, P2142 HCAPLUS
- (33) Esslinger, C; Human Gene Therapy 2002, V13, P1091 HCAPLUS
- (34) Evans, D; Immunological Reviews 2001, V183, P141 HCAPLUS
- (35) Fan, Z; Journal of Virology 2001, V75, P4413 HCAPLUS
- (36) Fanger, N; Journal of Immunology 1996, V157, P541 HCAPLUS
- (37) Fields, R; Proceedings of the National Academy of Sciences 1998, V95, P9482 HCAPLUS
- (38) Figdor, C; Nature Reviews Immunology 2002, V2, P77 HCAPLUS
- (39) Finke, J; Immunology Today 1999, V20, P158 HCAPLUS
- (40) Finzi, D; Nature Medicine 1999, V5, P512 HCAPLUS
- (41) Flamand, V; European Journal of Immunology 1994, V24, P605 MEDLINE
- (42) Foley, H; Journal of Virology 2002, V76, P19 HCAPLUS
- (43) Fong, L; Annual Review of Immunology 2000, V18, P245 HCAPLUS
- (44) Fong, L; Journal of Immunology 2001, V166, P4254 HCAPLUS
- (45) Fong, L; Proceedings of the American Association for Cancer Research 1999, V40, P85
- (46) Fong, L; Proceedings of the National Academy of Sciences 2001, V98, P8809 HCAPLUS
- (47) Gabrilovich, D; Nature Medicine 1996, V2, P1096 HCAPLUS
- (48) Galeà-Lauri, J; Cancer Immunology and Immunotherapy 2002, V51, P299 HCAPLUS
- (49) Gandhi, R; Annual Review of Medicine 2002, V53, P149 HCAPLUS
- (50) Geiger, J; Lancet 2000, V356, P1163 MEDLINE
- (51) Geijtenbeek, T; Cell 2000, V100, P587 HCAPLUS
- (52) Geijtenbeek, T; Journal of Leukocyte Biology 2002, V71, P921 HCAPLUS
- (53) Gilboa, E; Cancer Immunology and Immunotherapy 1998, V46, P82 HCAPLUS
- (54) Gnant, M; 7th International Symposium on Dendritic Cells 2002
- (55) Gong, J; Nature Medicine 1997, V3, P558 HCAPLUS
- (56) Grohmann, U; Journal of Immunology 1997, V158, P3593 HCAPLUS
- (57) Gruber, A; Blood 2000, V96, P1327 HCAPLUS
- (58) Guermontprez, P; Annual Review of Immunology 2002, V20, P621 HCAPLUS
- (59) Harrer, T; Journal of Immunology 1996, V156, P2616 HCAPLUS
- (60) Havlir, D; Journal of Infectious Diseases 2000, V182, P1658 MEDLINE
- (61) Hawiger, D; Journal of Experimental Medicine 2001, V194, P769 HCAPLUS
- (62) Heath, W; Annual Review of Immunology 2001, V19, P47 HCAPLUS
- (63) Heath, W; Nature Reviews Immunology 2001, V1, P126 HCAPLUS
- (64) Heiser, A; Cancer Research 2001, V61, P3388 HCAPLUS
- (65) Heiser, A; Journal of Clinical Investigation 2002, V109, P409 HCAPLUS
- (66) Holtl, L; Lancet 1998, V352, P1358 MEDLINE
- (67) Hsu, F; Nature Medicine 1996, V2, P52 HCAPLUS
- (68) Inaba, K; Journal of Experimental Medicine 1997, V186, P665 HCAPLUS

- (69) Inaba, K; Proceedings of the National Academy of Sciences 1983, V80, P6041
MEDLINE
- (70) Ingulli, E; Journal of Experimental Medicine 1997, V185, P2133 HCAPLUS
- (71) Jager, E; Current Opinion in Immunology 2002, V14, P178 HCAPLUS
- (72) Kalams, S; Journal of Virology 1999, V73, P6715 HCAPLUS
- (73) Kavanagh, D; Nature Immunology 2002, V3, P891 HCAPLUS
- (74) Kent, S; Immunological Reviews 2001, V183, P94 HCAPLUS
- (75) Klagge, I; Journal of General Virology 1999, V80, P823 HCAPLUS
- (76) Koch, F; Journal of Experimental Medicine 1996, V184, P741 HCAPLUS
- (77) Komanduri, K; Nature Medicine 1998, V4, P953 HCAPLUS
- (78) Kovacs, J; New England Journal of Medicine 2000, V342, P1416 HCAPLUS
- (79) Kundu, S; AIDS Research and Human Retroviruses 1998, V14, P551 MEDLINE
- (80) Lespagnard, L; International Journal of Cancer 1999, V84, P309 MEDLINE
- (81) Litzow, M; 7th International Symposium on Dendritic Cells 2002
- (82) Liu, K; Journal of Experimental Medicine 2002, V196, P1091 HCAPLUS
- (83) Lu, W; Nature Medicine 2003, V9, P27 HCAPLUS
- (84) Ludewig, B; Journal of Experimental Medicine 2000, V191, P795 HCAPLUS
- (85) Luster, A; Current Opinion in Immunology 2002, V14, P129 HCAPLUS
- (86) Lutz, M; Trends in Immunology 2002, V23, P445 HCAPLUS
- (87) Maier, T; 7th International Symposium on Dendritic Cells 2002
- (88) Marincola, F; Advances in Immunology 2000, V74, P181 HCAPLUS
- (89) Marovich, M; Journal of Infectious Diseases 2002, V185, P1242
- (90) Masurier, C; Journal of Virology 1998, P72
- (91) Matthes, C; 7th International Symposium on Dendritic Cells 2002
- (92) Mayordomo, J; Nature Medicine 1995, V1, P1297 HCAPLUS
- (93) McGrath, Y; 7th International Symposium on Dendritic Cells 2002
- (94) Migueles, S; Nature Immunology 2002, V3, P1061 HCAPLUS
- (95) Mitchell, D; Journal of Clinical Investigation 2000, V106, P1065 HCAPLUS
- (96) Mougneau, E; Journal of Experimental Medicine 2002, V196, P1013 HCAPLUS
- (97) Murphy, G; Prostate 1999, V38, P73 HCAPLUS
- (98) Nair, S; Nature Biotechnology 1998, V16, P364 HCAPLUS
- (99) Nehete, P; Vaccine 2001, V20, P813 HCAPLUS
- (100) Nestle, F; Nature Medicine 1998, V4, P328 HCAPLUS
- (101) Nestle, F; Nature Medicine 2001, V7, P761 HCAPLUS
- (102) Nimmerjahn, F; 7th International Symposium on Dendritic Cells 2002
- (103) Nishiyama, T; Clinical Cancer Research 2001, V7, P23 HCAPLUS
- (104) Nouri-Shirazi, M; Immunology Letters 2000, V74, P5 HCAPLUS
- (105) O'Connor, D; Immunological Reviews 2001, V183, P115 MEDLINE
- (106) Palella, F; New England Journal of Medicine 1998, V338, P853
- (107) Parmiani, G; Journal of the National Cancer Institute 2002, V94, P805
HCAPLUS
- (108) Paterson, D; Annals of Internal Medicine 2000, V133, P21 MEDLINE
- (109) Perumal, D; 9th Conference on Retroviruses and Opportunistic Infections
2002
- (110) Pohlmann, S; Trends in Immunology 2001, V22, P643 HCAPLUS
- (111) Prabakaran, I; Annals of Surgical Oncology 2002, V9, P411
- (112) Reddy, A; Blood 1997, V90, P3640 HCAPLUS
- (113) Reichardt, V; Blood 1999, V93, P2411 HCAPLUS
- (114) Reinhardt, P; 7th International Symposium on Dendritic Cells 2002
- (115) Romani, N; Journal of Immunological Methods 1996, V196, P137 HCAPLUS
- (116) Roncarolo, M; Journal of Experimental Medicine 2001, V193, P5 HCAPLUS
- (117) Rosenberg, E; Nature 2000, V407, P523 HCAPLUS
- (118) Rosenberg, E; Science 1997, V278, P1447 HCAPLUS
- (119) Saeki, H; Journal of Immunology 1999, V162, P2472 HCAPLUS
- (120) Salgaller, M; Cancer 1999, V86, P2674 MEDLINE
- (121) Sallusto, F; Journal of Experimental Medicine 1995, V182, P389 HCAPLUS
- (122) Sallusto, F; Journal of Experimental Medicine 1999, V189, P611 HCAPLUS
- (123) Santin, A; Journal of Virology 1999, V73, P5402 HCAPLUS
- (124) Santin, A; New England Journal of Medicine 2002, V346, P1752
- (125) Sastry, K; 7th International Symposium on Dendritic Cells 2002
- (126) Schadendorf, D; Recent Results in Cancer Research 2001, V158, P236
HCAPLUS
- (127) Schuler, G; Journal of Experimental Medicine 1997, V186, P1183 HCAPLUS

- (128) Schuler-Thurner, B; Journal of Experimental Medicine 2002, V195, P1279
HCAPLUS
- (129) Schwartz, D; AIDS Research and Human Retroviruses 1994, V10, P1703
MEDLINE
- (130) Sempowski, G; Annual Review of Medicine 2002, V53, P269 HCAPLUS
- (131) Shafran, S; Annals of Internal Medicine 2002, V137, P734
- (132) Shortman, K; Nature Reviews Immunology 2002, V2, P151 HCAPLUS
- (133) Small, E; Journal of Clinical Oncology 2000, V18, P3894 HCAPLUS
- (134) Smith, K; Journal of Infectious Diseases 2002, V185, PS123 HCAPLUS
- (135) Steinman, R; Cell 2000, V100, P491 HCAPLUS
- (136) Steinman, R; Journal of Clinical Investigation 2002, V109, P1519 HCAPLUS
- (137) Steinman, R; Journal of Experimental Medicine 1973, V137, P1142 MEDLINE
- (138) Steinman, R; Journal of Experimental Medicine 1974, V139, P380 MEDLINE
- (139) Steinman, R; Proceedings of the National Academy of Sciences 2002, V99,
P351 HCAPLUS
- (140) Sullenger, B; Nature 2002, V418, P252 HCAPLUS
- (141) Svensson, M; Journal of Immunology 1997, V158, P4229 HCAPLUS
- (142) Thurner, B; Journal of Experimental Medicine 1999, V190, P1669 HCAPLUS
- (143) Timmerman, J; Annual Review of Medicine 1999, V50, P507 HCAPLUS
- (144) Tjoa, B; Prostate 1999, V40, P125 MEDLINE
- (145) Triozzi, P; Cancer 2000, V89, P2646 MEDLINE
- (146) Turley, S; Science 2000, V288, P522 HCAPLUS
- (147) Tuting, T; Journal of Immunology 1998, V160, P1139 HCAPLUS
- (148) van Baalen, C; Trends in Immunology 2002, V23, P478 HCAPLUS
- (149) Vicari, A; Seminars in Cancer Biology 2002, V12, P33 HCAPLUS
- (150) Wagner, G; AIDS Research and Human Retroviruses 2002, V18, P143
- (151) Weissman, D; Journal of Immunology 2000, V165, P4710 HCAPLUS
- (152) Weissman, D; Proceedings of the National Academy of Sciences 1995, V92,
P826 HCAPLUS
- (153) Wilson, C; Journal of Immunology 1999, V162, P3070 HCAPLUS
- (154) Yi, Q; British Journal of Haematology 2001, V117, P297
- (155) Zhang, J; World Journal of Gastroenterology 2001, V7, P690 HCAPLUS
- (156) Zhao, X; Journal of Virology 2002, V76, P3007 HCAPLUS
- (157) Zhu, Y; 9th Conference on Retroviruses and Opportunistic Infections 2002
- (158) Zinkernagel, R; Seminars in Immunology 2000, V12, P163 HCAPLUS
- (159) Zitvogel, L; Journal of Experimental Medicine 1996, V183, P87 HCAPLUS

L40 ANSWER 6 OF 15 HCAPLUS COPYRIGHT 2004 ACS on STN

AN 2003:2999 HCAPLUS

DN 138:121297

ED Entered STN: 03 Jan 2003

TI Therapeutic **dendritic-cell** vaccine for simian AIDS

AU (Lu, Wei; Wu, Xiaoxian; Lu, Yaozeng; Guo, Weizhong; **Andrieu, Jean-Marie**

CS. Institut de Recherche sur les Vaccins et l'Immunotherapie des Cancers et du Sida, Paris, Fr.

SO Nature Medicine (New York, NY, United States) (2003), 9(1), 27-32
CODEN: NAMEFI; ISSN: 1078-8956

PB Nature Publishing Group

DT Journal

LA English

CC 15-2 (Immunochemistry)

AB An effective immune response against **human**

immunodeficiency virus or simian immunodeficiency virus

(SIV) is critical in achieving control of viral replication. Here, the authors show in SIV-infected rhesus monkeys that an effective and durable SIV-specific cellular and humoral immunity is elicited by a vaccination with chemical **inactivated SIV-pulsed dendritic cells**. After 3 immunizations made at 2-wk intervals, the animals exhibited a 50-fold decrease of SIV DNA and a 1000-fold decrease of SIV RNA in peripheral blood. Such reduced viral load levels were maintained over the remaining 34 wk of the study. Mol. and cellular analyses of axillary and inguinal node lymphocytes of vaccinated monkeys revealed a

correlation between decreased SIV DNA and RNA levels and increased SIV-specific T-cell responses. Neutralizing antibody responses were augmented and remained elevated. **Inactivated whole virus-pulsed dendritic cell** vaccines are promising means to control diseases caused by immuno- deficiency viruses.

- ST **dendritic cell inactivated virus vaccine**
simian AIDS
- IT Vaccines
(AIDS; **inactivated whole virus-pulsed dendritic cell vaccine** for simian AIDS)
- IT T cell (lymphocyte)
(**activation; inactivated whole virus-pulsed dendritic cell vaccine** for simian AIDS)
- IT Adoptive immunotherapy
Dendritic cell
Macaca mulatta
Simian immunodeficiency virus
(**inactivated whole virus-pulsed dendritic cell vaccine** for simian AIDS)
- IT Antibodies
RL: BSU (Biological study, unclassified); BIOL (Biological study)
(neutralizing; **inactivated whole virus-pulsed dendritic cell vaccine** for simian AIDS)
- IT Anti-AIDS agents
(vaccines; **inactivated whole virus-pulsed dendritic cell vaccine** for simian AIDS)

RE.CNT 37 THERE ARE 37 CITED REFERENCES AVAILABLE FOR THIS RECORD
RE

- (1) Andrieu, J; Immunol Today 1995, V16, P5 HCAPLUS
- (2) Barouch, D; Nature 2002, V415, P335 HCAPLUS
- (3) Buseyne, F; Nature Med 2001, V7, P344 HCAPLUS
- (4) Champagne, P; Nature 2001, V410, P106 HCAPLUS
- (5) Chougnet, C; J Immunol 1999, V163, P1666 HCAPLUS
- (6) Cohen, O; Immunol Rev 1997, V159, P31 HCAPLUS
- (7) Davey, R; Proc Natl Acad Sci USA 1999, V96, P15109 HCAPLUS
- (8) Donaghy, H; Blood 2001, V98, P2574 HCAPLUS
- (9) Feldman, S; Clin Immunol 2001, V101, P201 HCAPLUS
- (10) Fenyo, E; Immunol Lett 1996, V51, P95 MEDLINE
- (11) Grabar, S; Ann Intern Med 2000, V133, P401 HCAPLUS
- (12) Grassi, F; Aids 1999, V13, P759 MEDLINE
- (13) Gray, C; J Immunol 1999, V162, P1780 HCAPLUS
- (14) Hatano, H; Aids 2000, V14, P1357 HCAPLUS
- (15) Hermans, I; J Immunol 2000, V164, P3095 HCAPLUS
- (16) Hirsch, V; Adv Pharmacol 2000, V49, P437 MEDLINE
- (17) Jin, X; Mol Med 2000, V6, P803 HCAPLUS
- (18) Kalams, S; J Virol 1999, V73, P6721 HCAPLUS
- (19) Knight, S; Curr Opin Immunol 1993, V5, P374 MEDLINE
- (20) Kostense, S; Blood 2002, V99, P2505 HCAPLUS
- (21) Lederman, M; JAMA 2000, V284, P223 MEDLINE
- (22) Lieberman, J; Blood 2001, V98, P1667 HCAPLUS
- (23) Lu, W; Adv Exp Med Biol 1995, V374, P235 HCAPLUS
- (24) Lu, W; Blood 2000, V96, P250 HCAPLUS
- (25) Lu, W; J Immunol 2001, V167, P2929 HCAPLUS
- (26) Lu, W; J Virol 2001, V75, P8949 HCAPLUS
- (27) Lu, W; Nature Med 1999, V5, P1081 HCAPLUS
- (28) Ludewig, B; J Immunol 1999, V163, P1839 HCAPLUS
- (29) McIlroy, D; AIDS Res Hum Retroviruses 1998, V14, P505 HCAPLUS
- (30) Mehlhop, E; J Immunol Methods 2002, V260, P219 HCAPLUS
- (31) Nair, S; Nature Biotechnol 1998, V16, P364 HCAPLUS
- (32) Norbury, C; Nature Immunol 2002, V3, P265 HCAPLUS
- (33) Pacanowski, J; Blood 2001, V98, P3016 HCAPLUS
- (34) Rossio, J; J Virol 1998, V72, P7992 HCAPLUS
- (35) Salerno-Goncalves, R; J Virol 2000, V74, P6648 HCAPLUS

- (36) Sapp, M; Immunol Lett 1999, V66, P121 HCAPLUS
 (37) Zimmer, M; Blood 2002, V99, P2859 HCAPLUS

L40 ANSWER 7 OF 15 HCAPLUS COPYRIGHT 2004 ACS on STN

AN 2002:220375 HCAPLUS

DN 136:241666

ED Entered STN: 22 Mar 2002

TI Means for regulating immune defenses

IN **Andrieu, Jean-Marie**; Lu, Wei; Achour, Amar

PA Institut Necker, Fr.

SO PCT Int. Appl., 39 pp.

CODEN: PIXXD2

DT Patent

LA French

IC ICM A61K031-5415

ICS A61P031-00; A61P031-06; A61P031-18; A61P035-00

CC 1-7 (Pharmacology)

Section cross-reference(s): 15, 63

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2002022130	A2	20020321	WO 2001-FR2810	20010911
	WO 2002022130	A3	20020613		
	W:				
	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,				
	CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,				
	GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,				
	LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL,				
	PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG,				
	US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
	RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY,				
	DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF,				
	BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	FR 2813792	A1	20020315	FR 2000-11567	20000912
	FR 2813792	B1	20030314		
	AU 2001091934	A5	20020326	AU 2001-91934	20010911
PRAI	FR 2000-11567	A	20000912		
	WO 2001-FR2810	W	20010911		

OS MARPAT 136:241666

AB The invention concerns the rebuilding of immune defenses. More particularly, it concerns the use of a compound capable of acting on monocytes/macrophages and/or **dendritic cells**, so as to provide them with a signal inducing proliferation of lymphocytes, in particular T lymphocytes, providing thereby a pos. regulation of immune defenses in the treated organism. The **active** substance is advantageously selected among phenazine, phenoxazine and phenothiazine derivs. in particular, aminoperazine [2-amino-10-[3'-(1-methyl-4-piperazinyl)-propyl]phenothiazine]. Among the examples provided are effects of aminoperazine on immune function (T cell survival, proliferation and antiviral **activity**, cytokine production) in vitro and in HIV-pos. patients. Such derivs. may also be useful in treating tumors, infections and other immune deficiencies.

ST aminoperazine HIV1 T lymphocyte **dendritic cell**; immune deficiency phenothiazine deriv monocyte macrophage; antitumor infection immunity T cell aminoperazine

IT Antiviral agents

(antiviral **activity** of CD8-pos. T cells sensitized by HIV p24gag)

IT Drug delivery systems

(injections; phenothiazine derivs. for rebuilding immune function via T cell proliferation)

IT Drug delivery systems

(lozenges; phenothiazine derivs. for rebuilding immune function via T cell proliferation)

- IT Antitumor agents
(metastasis; aminoperazine for rebuilding immune function via T cell proliferation)
- IT gag proteins
RL: BSU (Biological study, unclassified); BIOL (Biological study)
(p24gag; antiviral activity of CD8-pos. T cells sensitized by HIV p24gag)
- IT Anti-AIDS agents
Antitumor agents
Apoptosis
CD4-positive T cell
CD8-positive T cell
Dendritic cell
Human
Human immunodeficiency virus 1
Immunodeficiency
Infection
Macrophage
Monocyte
T cell (lymphocyte)
(phenothiazine derivs. for rebuilding immune function via T cell proliferation)
- IT Cytokines
Ki-67 antigen
RL: BSU (Biological study, unclassified); BIOL (Biological study)
(phenothiazine derivs. for rebuilding immune function via T cell proliferation)
- IT Interleukin 1 receptor antagonist
Interleukin 10
Interleukin 12
Interleukin 15
Interleukin 18
Interleukin 1 β
Interleukin 4
Interleukin 6
Tumor necrosis factors
RL: BSU (Biological study, unclassified); BIOL (Biological study)
(phenothiazine derivs. for rebuilding immune function via T cell proliferation: effect on cytokine production)
- IT Viral DNA
Viral RNA
RL: BSU (Biological study, unclassified); BIOL (Biological study)
(phenothiazine derivs. for rebuilding immune function via T cell proliferation: effect on proviral DNA and viral RNA in HIV -pos. patients)
- IT T cell (lymphocyte)
(proliferation; phenothiazine derivs. for rebuilding immune function via T cell proliferation)
- IT Drug delivery systems
(solns.; phenothiazine derivs. for rebuilding immune function via T cell proliferation)
- IT Drug delivery systems
(suspensions; phenothiazine derivs. for rebuilding immune function via T cell proliferation)
- IT Drug delivery systems
(tablets; phenothiazine derivs. for rebuilding immune function via T cell proliferation)
- IT Immunization
(vaccination; phenothiazine derivs. for rebuilding immune function via T cell proliferation combined with vaccines)
- IT Transforming growth factors
RL: BSU (Biological study, unclassified); BIOL (Biological study)
(β 1-; phenothiazine derivs. for rebuilding immune function via T

cell proliferation: effect on cytokine production)

IT Interferons
 RL: BSU (Biological study, unclassified); BIOL (Biological study)
 (γ; phenothiazine derivs. for rebuilding immune function via T
 cell proliferation: effect on cytokine production)

IT 92-82-0D, Phenazine, derivs. 92-84-2D, Phenothiazine, derivs.
 135-67-1D, Phenoxazine, derivs.
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (aminoperazine for rebuilding immune function via T cell proliferation)

IT 50-53-3, Chlorpromazine, biological studies
 RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL
 (Biological study); USES (Uses)
 (aminoperazine for rebuilding immune function via T cell proliferation:
 comparisons with other phenothiazines)

IT 367274-46-2
 RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL
 (Biological study); USES (Uses)
 (phenothiazine derivs. for rebuilding immune function via T cell
 proliferation)

IT 117-89-5, Trifluoperazine 3937-85-7 107457-56-7 404825-16-7
 404825-17-8 404825-18-9 404825-19-0 404825-20-3 404825-21-4
 404825-22-5 404825-23-6 404825-24-7
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (phenothiazine derivs. for rebuilding immune function via T cell
 proliferation)

L40 ANSWER 8 OF 15 HCAPLUS COPYRIGHT 2004 ACS on STN

AN 2001:705874 HCAPLUS

DN 136:4662

ED Entered STN: 27 Sep 2001

TI In vitro **human immunodeficiency virus**
 eradication by autologous CD8+ T cells expanded with
inactivated-virus-pulsed-dendritic
cells

AU Lu, Wei; Andrieu, Jean-Marie

CS Laboratory of Molecular Oncology and Virology, Necker Faculty of Medicine
 at Saints-Peres Biomedical Center, Rene Descartes University, Paris,
 75270, Fr.

SO Journal of Virology (2001), 75(19), 8949-8956

CODEN: JOVIAM; ISSN: 0022-538X

PB American Society for Microbiology

DT Journal

LA English

CC 15-10 (Immunochemistry)

Section cross-reference(s): 1

AB Despite significant immune recovery with potent highly **active**
 antiretroviral therapy (HAART), eradication of **human**
immunodeficiency virus (HIV) from the bodies
 of infected individuals represents a challenge. The authors hypothesized
 that an inadequate or inappropriate signal in virus-specific antigen
 presentation might contribute to the persistent failure to mount efficient
 anti-HIV immunity in most HIV-infected individuals.
 Here, they conducted an in vitro study with untreated and HAART-treated
 HIV type 1 (HIV-1) patients which showed that
pulsing of monocyte-derived **dendritic cells**
 (DC) with aldrithiol-2-**inactivated** autologous virus resulted in
 the expansion of virus-specific CD8+ T cells which were capable
 of killing HIV-1-infected cells and eradicating the virus from
 cultured patient peripheral blood mononuclear cells independently of the
 disease stages and HAART response statuses of the patients. This in vitro
 anti-HIV effect was further enhanced by the HIV
 protease inhibitor indinavir (at a nonantiviral concentration), which has been
 shown previously to be able to up-regulate directly patient T-cell

Applicant

proliferation following immune stimulation. However, following a 2-day treatment with culture supernatant derived from immune-activated T cells (which mimics an in vivo environment of HIV-disseminated and immune-activated lymphoid tissues), DC lost their capacity to present de novo inactivated virus-derived antigens. These findings provide important information for understanding the establishment of chronic HIV infection and indicate a perspective for clinical use of DC-based therapeutic vaccines against HIV.

- ST HIV CD8 T cell virus pulsed dendritic cell HAART
- IT Anti-AIDS agents
 - CD8-positive T cell
 - Dendritic cell
 - Human immunodeficiency virus 1
 - (HIV-1 eradication by autologous CD8+ T cells expanded with inactivated virus-pulsed dendritic cells)
- IT Antigen presentation
 - (HIV-1 eradication by autologous CD8+ T cells expanded with inactivated virus-pulsed dendritic cells in relation to)
- IT Anti-AIDS agents
 - (vaccines; HIV-1 eradication by autologous CD8+ T cells expanded with inactivated virus-pulsed dendritic cells)
- IT 150378-17-9, Indinavir
 - RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 - (HIV-1 eradication by autologous CD8+ T cells expanded with inactivated virus-pulsed dendritic cells in presence of)

RE.CNT 30 THERE ARE 30 CITED REFERENCES AVAILABLE FOR THIS RECORD
RE

- (1) Bohler, T; Blood 2001, V97, P1898 HCAPLUS
- (2) Cavert, W; Science 1997, V276, P960 HCAPLUS
- (3) Chun, T; Nat Med 1999, V5, P651 HCAPLUS
- (4) Chun, T; Proc Natl Acad Sci USA 1997, V94, P13193 HCAPLUS
- (5) Collier, A; N Engl J Med 1996, V334, P1011 HCAPLUS
- (6) Davey, R; Proc Natl Acad Sci USA 1999, V96, P15109 HCAPLUS
- (7) Finzi, D; Nat Med 1999, V5, P512 HCAPLUS
- (8) Gharar, S; Ann Intern Med 2000, V133, P401
- (9) Gray, C; J Immunol 1999, V162, P1780 HCAPLUS
- (10) Harrigan, P; AIDS 1999, V13, PF59 HCAPLUS
- (11) Hatanó, H; AIDS 2000, V14, P1357 HCAPLUS
- (12) Haynes, B; Science 1996, V271, P324 HCAPLUS
- (13) Ibanez, A; AIDS 1999, V13, P1045 HCAPLUS
- (14) Kalams, S; J Virol 1999, V73, P6721 HCAPLUS
- (15) Kaufmann, D; Lancet 1998, V351, P723 MEDLINE
- (16) Ledergerber, B; Lancet 1999, V353, P863 MEDLINE
- (17) Levitz, S; N Engl J Med 1998, V338, P1074 MEDLINE
- (18) Lu, W; Adv Exp Med Biol 1995, V374, P235 HCAPLUS
- (19) Lu, W; Blood 2000, V96, P250 HCAPLUS
- (20) Lu, W; Blood 2001, V97, P1900 HCAPLUS
- (21) Lu, W; Nat Med 1999, V5, P1081 HCAPLUS
- (22) Lu, W; Nature 1994, V368, P269 MEDLINE
- (23) McMichael, A; Annu Rev Immunol 1997, V15, P271 HCAPLUS
- (24) Rossio, J; J Virol 1998, V72, P7992 HCAPLUS
- (25) Salerno-Goncalves, R; Immunol Lett 1998, V64, P71 HCAPLUS
- (26) Salerno-Goncalves, R; J Virol 2000, V74, P6648 HCAPLUS
- (27) Santini, S; J Exp Med 2000, V191, P1777 HCAPLUS
- (28) Sloand, E; Blood 1999, V94, P1021 HCAPLUS
- (29) Wilson, C; J Immunol 1999, V162, P3070 HCAPLUS
- (30) Zhang, Z; Science 1999, V286, P1353 HCAPLUS

L40 ANSWER 9 OF 15 HCAPLUS COPYRIGHT 2004 ACS on STN
AN 2001:630088 HCAPLUS
DN 135:302780
ED Entered STN: 30 Aug 2001
TI Enhanced **dendritic cell**-driven proliferation and anti-
HIV activity of CD8+ T cells by a new phenothiazine
derivative, aminoperazine
AU Lu, Wei; Achour, Amar; Arlie, Marine; Cao, Li; **Andrieu, Jean-Marie**
CS Laboratory of Molecular Oncology and Virology, Necker Faculty of Medicine,
Saints-Peres Biomedical Center, Rene Descartes University, Paris, 75270,
Fr.
SO Journal of Immunology (2001), 167(5), 2929-2935
CODEN: JOIMA3; ISSN: 0022-1767
PB American Association of Immunologists
DT Journal
LA English
CC 15-8 (Immunochemistry)
AB T cell anergy, apoptosis, and chronic **activation** of T
lymphocytes are prevailing features of **HIV** infection. The
inability to develop an efficient natural antiviral **activity** in
infected patients might be the consequence of a failure of the Ag
presentation by **dendritic cells** (DCs) in chronically
activated lymphoid tissues. We have identified a new
phenothiazine derivative aminoperazine (APR; 2-amino-10-[3'-(1-methyl-4-
piperazinyl)propyl]phenothiazine, C₂₀H₂₆N₄S; m.w. 354.51) able to increase
(ED from 0.1 to 100 nM) the Ag-specific DC-driven proliferation and
differentiation of in vitro **HIV**-infected and uninfected normal
donor T cells and of T cells from **HIV**-1-infected patients. The
immunomodulatory effect of APR-sensitized DCs were ascribed to soluble
factors derived from DCs. APR was also capable of increasing **HIV**
gag-p24-specific proliferation and anti-**HIV** cytotoxic
activity of patients' CD8+ T cells against autologous
B-lymphoblastoid cell lines expressing a **HIV** gag gene, resulting
in the suppression of both proviral DNA and supernatant viral RNA in the
HIV-1-infected patients' T cell culture. This new phenothiazine
derivative (APR) might be used for boosting the immune response of vaccinated
individuals and for restoring the immunity of immunocompromised patients.
ST **HIV**1 gag protein T lymphocyte aminoperazine
IT Immune tolerance
(anergy; phenothiazine derivative, aminoperazine in anti-**HIV**
activity of CD8+ T cells)
IT gag proteins
RL: BAC (Biological activity or effector, except adverse); BSU (Biological
study, unclassified); BIOL (Biological study)
(p24gag; phenothiazine derivative, aminoperazine in anti-**HIV**
activity of CD8+ T cells)
IT Apoptosis
CD8-positive T cell
Cytotoxicity
Human immunodeficiency virus 1
(phenothiazine derivative, aminoperazine in anti-**HIV**
activity of CD8+ T cells)
IT Interleukin 10
Interleukin 12
Interleukin 15
Interleukin 18
Interleukin 1 β
Interleukin 4
Interleukin 6
Lymphotoxin
Tumor necrosis factors
RL: BOC (Biological occurrence); BSU (Biological study, unclassified);

BIOL (Biological study); OCCU (Occurrence)
 (phenothiazine derivative, aminoperazine in anti-HIV
activity of CD8+ T cells and the expression of)

IT Interferons
 RL: BOC (Biological occurrence); BSU (Biological study, unclassified);
 BIOL (Biological study); OCCU (Occurrence)
 (γ; phenothiazine derivative, aminoperazine in anti- HIV
activity of CD8+ T cells and the expression of)

IT 367274-46-2
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological
 study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES
 (Uses)
 (phenothiazine derivative, aminoperazine in anti-HIV
activity of CD8+ T cells)

IT 92-84-2D, Phenothiazine, derivs.
 RL: BSU (Biological study, unclassified); BIOL (Biological study)
 (phenothiazine derivative, aminoperazine in anti-HIV
activity of CD8+ T cells)

RE.CNT 35 THERE ARE 35 CITED REFERENCES AVAILABLE FOR THIS RECORD
 RE

- (1) Andersson, J; AIDS 1998, V12, PF123 HCAPLUS
- (2) Andrieu, J; Immunol Today 1995, V16, P5 HCAPLUS
- (3) Badley, A; J Exp Med 1997, V185, P55 MEDLINE
- (4) Brossart, P; Blood 1997, V90, P1594 HCAPLUS
- (5) Cavert, W; Science 1997, V276, P960 HCAPLUS
- (6) Chun, T; Proc Natl Acad Sci 1997, V94, P13193 HCAPLUS
- (7) Collier, A; N Engl J Med 1996, V334, P1011 HCAPLUS
- (8) Fauci, A; Science 1993, V262, P1011 HCAPLUS
- (9) Finkel, T; Nat Med 1995, V1, P129 HCAPLUS
- (10) Finzi, D; Nat Med 1999, V5, P512 HCAPLUS
- (11) Fleury, S; Nat Med 1998, V4, P794 HCAPLUS
- (12) Fleury, S; Proc Natl Acad Sci 2000, V97, P5393 HCAPLUS
- (13) Grabar, S; Ann Intern Med 2000, V133, P401 HCAPLUS
- (14) Gray, C; J Immunol 1999, V162, P1780 HCAPLUS
- (15) Hellerstein, M; Nat Med 1999, V5, P83 HCAPLUS
- (16) Hosmalin, A; J Immunol 1992, V149, P1311 HCAPLUS
- (17) Ibanez, A; AIDS 1999, V13, P1045 HCAPLUS
- (18) Kalams, S; J Virol 1999, V73, P6721 HCAPLUS
- (19) Kaneko, H; Clin Exp Immunol 1997, V109, P41 HCAPLUS
- (20) Kaufmann, D; Lancet 1998, V351, P723 MEDLINE
- (21) Knight, S; AIDS 1996, V10, P807 MEDLINE
- (22) Ledergerber, B; Lancet 1999, V353, P863 MEDLINE
- (23) Levitz, S; N Engl J Med 1998, V338, P1074 MEDLINE
- (24) Lu, L; J Immunol 1997, V158, P5676 HCAPLUS
- (25) Lu, W; Blood 2000, V96, P250 HCAPLUS
- (26) Lu, W; Blood 2001, V97, P1900 HCAPLUS
- (27) Lu, W; Nat Med 1999, V5, P1081 HCAPLUS
- (28) Lu, W; Nature 1994, V368, P269 MEDLINE
- (29) Miedema, F; Immunodef Rev 1992, V3, P173 MEDLINE
- (30) Nelson, D; J Immunol 2000, V165, P6123 HCAPLUS
- (31) Salerno-Goncalves, R; Immunol Lett 1998, V64, P71 HCAPLUS
- (32) Salerno-Goncalves, R; J Virol 2000, V74, P6648 HCAPLUS
- (33) Schols, D; J Virol 1996, V70, P4953 HCAPLUS
- (34) Sloand, E; Blood 1999, V94, P1021 HCAPLUS
- (35) Spinillo, A; Gynecol Oncol 1993, V48, P210 MEDLINE

L40 ANSWER 10 OF 15 HCAPLUS COPYRIGHT 2004 ACS on STN

AN 1998:620226 HCAPLUS

ED Entered STN: 01 Oct 1998

TI **Dendritic cells route human
 immunodeficiency virus** to lymph nodes after vaginal or
 intravenous administration to mice

AU Masurier, Carole; Salomon, Benoit; Guettari, Nadia; Pioche, Catherine;

CS Lachapelle, Francois; Guigon, Martine; Klatzmann, David
 Laboratoire de Biologie et Therapeutique des Pathologies Immunitaires,
 Universite Pierre et Marie Curie/CNRS ESA 70-87, Hopital
 Pitie-Salpetriere, Paris, 75651, Fr.

SO Journal of Virology (1998), 72(10), 7822-7829
 CODEN: JOVIAM; ISSN: 0022-538X

PB American Society for Microbiology

DT Journal

LA English

AB We have developed a murine model to study the involvement of
dendritic cells (DC) in human immunodeficiency virus (HIV) routing from an
 inoculation site to the lymph nodes (LN). Murine bone marrow-derived DC
 migrate to the draining LN within 24 h after s.c. injection. After
 incubation of these cells with heat-inactivated (Hi) HIV
 type 1 (HIV-1), HIV RNA sequences were detected in the
 draining LN only. Upon injection of DC **pulsed** with infectious
 HIV, the virus recovered in the draining LN was still able to
 productively infect human T cells. After a vaginal challenge with Hi
 HIV-1, the virus could be detected in the iliac and sacral
 draining LN at 24 h after injection. After an i.v. challenge, the virus
 could be detected in peripheral LN as soon as 30 min after injection. The
 specific depletion of a myeloid-related LN DC population, previously shown
 to take up blood macromols. and to translocate them into the LN, prevented
 HIV transport to LN. Together, our data demonstrate the critical
 role of DC for HIV routing to LN after either a vaginal or an
 i.v. challenge, which does not require their infection. Therefore, despite
 the fact that the mouse is not infectable by HIV, this small
 animal model might be useful to test preventive strategies against
 HIV.

RE.CNT 56 THERE ARE 56 CITED REFERENCES AVAILABLE FOR THIS RECORD
 RE

- (1) Allaerts, W; J Neuroimmunol 1997, V78, P184 HCAPLUS
- (2) Austyn, J; J Exp Med 1988, V167, P646 MEDLINE
- (3) Austyn, J; J Exp Med 1996, V183, P1287 HCAPLUS
- (4) Balter, M; Science 1996, V274, P1464 HCAPLUS
- (5) Barrat-Boyes, S; J Immunol 1997, V158, P4543
- (6) Blauvelt, A; J Clin Invest 1997, V100, P2043 HCAPLUS
- (7) Browning, J; Proc Natl Acad Sci USA 1997, V94, P14637 HCAPLUS
- (8) Cameron, P; AIDS Res Hum Retroviruses 1994, V10, P61 MEDLINE
- (9) Cameron, P; J Leukoc Biol 1994, V56, P257 HCAPLUS
- (10) Cameron, P; J Leukoc Biol 1996, V59, P158 HCAPLUS
- (11) Cameron, P; Science 1992, V257, P383 MEDLINE
- (12) Chakrabarti, L; Am J Pathol 1994, V144, P1226 MEDLINE
- (13) Clapham, P; Virology 1987, V158, P44 HCAPLUS
- (14) Connor, R; AIDS Res Hum Retroviruses 1994, V10, P321 MEDLINE
- (15) Embretson, J; Nature 1993, V362, P359 MEDLINE
- (16) Fossum, S; Scand J Immunol 1988, V27, P97 MEDLINE
- (17) Guery, J; J Exp Med 1996, V183, P751 HCAPLUS
- (18) Hill, S; Immunology 1990, V71, P277 MEDLINE
- (19) Ibrahim, M; Immunol Today 1995, V16, P181 HCAPLUS
- (20) Inaba, K; J Exp Med 1992, V175, P1157 HCAPLUS
- (21) Inaba, K; J Exp Med 1992, V176, P1693 MEDLINE
- (22) Klatzmann, D; Science 1984, V225, P59 MEDLINE
- (23) Knight, S; AIDS 1996, V10, P807 MEDLINE
- (24) Knight, S; Cell Immunol 1985, V94, P435 HCAPLUS
- (25) Knight, S; J Leukoc Biol 1997, V62, P78 HCAPLUS
- (26) Kupiec-Weglinski, J; J Exp Med 1988, V167, P632 MEDLINE
- (27) Lambrecht, B; J Immunol 1998, V160, P4090 HCAPLUS
- (28) Leenen, P; J Immunol 1998, V160, P2166 HCAPLUS
- (29) Leenen, P; Personal communication
- (30) Lores, P; AIDS Res Hum Retroviruses 1992, V8, P2063 MEDLINE
- (31) Lu, L; Transplantation 1995, V60, P1539 HCAPLUS

- (32) Macatonia, S; Immunology 1986, V59, P509 HCAPLUS
- (33) Macatonia, S; J Exp Med 1987, V166, P1654 MEDLINE
- (34) Marx, P; Nat Med 1996, V10, P1084
- (35) McWilliam, A; J Exp Med 1994, V179, P1331 MEDLINE
- (36) Metlay, J; J Exp Med 1990, V171, P1753 HCAPLUS
- (37) Miller, C; Lab Invest 1992, V68, P129
- (38) Morikawa, Y; Immunology 1995, V85, P575 HCAPLUS
- (39) Pantaleo, G; Nature 1993, V352, P355
- (40) Parr, M; Biol Reprod 1991, V45, P261 MEDLINE
- (41) Parr, M; J Reprod Immunol 1990, V17, P101 MEDLINE
- (42) Parr, M; Lab Invest 1994, V70, P369 MEDLINE
- (43) Pope, M; AIDS Res Hum Retroviruses 1997, V13, P819 MEDLINE
- (44) Pope, M; Cell 1994, V78, P389 MEDLINE
- (45) Roy, S; Immunology 1988, V64, P233 HCAPLUS
- (46) Sallusto, F; J Exp Med 1995, V182, P389 HCAPLUS
- (47) Salomon, B; Adv Exp Med Biol 1995, V378, P485 MEDLINE
- (48) Salomon, B; J Immunol 1994, V152, P537 MEDLINE
- (49) Salomon, B; J Immunol 1998, V160, P708 MEDLINE
- (50) Soto-Ramirez, L; Science 1996, V271, P1291 HCAPLUS
- (51) Spira, A; J Exp Med 1996, V183, P215 HCAPLUS
- (52) Steinman, R; Annu Rev Immunol 1991, V9, P271 HCAPLUS
- (53) Wain-Hobson, S; Science 1991, V252, P961 HCAPLUS
- (54) Weissman, D; Clin Microbiol Rev 1997, V10, P358 MEDLINE
- (55) Witmer-Pack, M; J Cell Sci 1993, V105, P965
- (56) Zack, J; Cell 1990, V61, P213 MEDLINE

L40 ANSWER 11 OF 15 HCAPLUS COPYRIGHT 2004 ACS on STN

AN 1997:503019 HCAPLUS

DN 127:107988

ED Entered STN: 09 Aug 1997

TI Methods for in vivo T cell **activation** by antigen-pulsed dendritic cells

IN Engleman, Edgar G.; Levy, Ronald; Hsu, Frank; Benike, Claudia

PA Board of Trustees of the Leland Stanford Junior University, USA

SO PCT Int. Appl., 48 pp.

CODEN: PIXXD2

DT Patent

LA English

IC ICM A61K035-14

ICS A61K039-02; A61K039-12; A61K039-385; A61K039-395; C12N005-08; G01N033-554

CC 15-2 (Immunochemistry)

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9722349	A1	19970626	WO 1996-US19954	19961217
	W: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, FI, GE, HU, IL, IS, JP, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TM, TR, TT, UA, UZ, VN, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM RW: KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
	AU 9711523	A1	19970714	AU 1997-11523	19961217
PRAI	US 1995-575432		19951220		
	WO 1996-US19954		19961217		

AB The present invention relates to methods of using isolated human dendritic cells to present exogenous antigens for the induction of immune responses in vivo. In particular, it relates to the isolation of dendritic cells from human blood, exposing the cells to lymphoma-derived Igs or to proteins derived from human immunodeficiency virus (HIV) as antigens, and infusing the antigen-pulsed dendritic cells into patients to induce and/or augment an antigen-specific immune response. The methods of the invention

described herein have a wide range of applications, including, but not limited to, the clin. use of antigen-pulsed dendritic cells as vaccines and/or immunotherapeutics against cancer and infectious agents such as viruses.

- ST virucide dendritic cell T cell **activation**; immunostimulant
dendritic cell T cell **activation**
- IT B cell (lymphocyte)
T cell (lymphocyte)
(-mediated immune response; in vivo T cell **activation** by
antigen-pulsed dendritic cells)
- IT Proteins, specific or class
RL: BAC (Biological activity or effector, except adverse); BPR (Biological
process); BSU (Biological study, unclassified); THU (Therapeutic use);
BIOL (Biological study); PROC (Process); USES (Uses)
(antigenic; in vivo T cell **activation** by antigen-pulsed
dendritic cells)
- IT Virus
(as antigen; in vivo T cell **activation** by antigen-pulsed
dendritic cells)
- IT Immunoglobulins
RL: BAC (Biological activity or effector, except adverse); BPR (Biological
process); BSU (Biological study, unclassified); THU (Therapeutic use);
BIOL (Biological study); PROC (Process); USES (Uses)
(as antigens; in vivo T cell **activation** by antigen-pulsed
dendritic cells)
- IT Blood
(dendritic cell isolation from human; in vivo T cell **activation**
by antigen-pulsed dendritic cells)
- IT Human immunodeficiency virus
(gp160 of; in vivo T cell **activation** by antigen-pulsed
dendritic cells)
- IT Envelope proteins
RL: BAC (Biological activity or effector, except adverse); BPR (Biological
process); BSU (Biological study, unclassified); THU (Therapeutic use);
BIOL (Biological study); PROC (Process); USES (Uses)
(gp160env, as antigen; in vivo T cell **activation** by
antigen-pulsed dendritic cells)
- IT Dendritic cell
Immunostimulants
(in vivo T cell **activation** by antigen-pulsed dendritic cells)
- IT Antigens
RL: BAC (Biological activity or effector, except adverse); BSU (Biological
study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES
(Uses)
(in vivo T cell **activation** by antigen-pulsed dendritic cells)
- IT Antitumor agents
(lymphoma; in vivo T cell **activation** by antigen-pulsed
dendritic cells)
- IT Antigens
RL: BAC (Biological activity or effector, except adverse); BSU (Biological
study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES
(Uses)
(tumor-associated; in vivo T cell **activation** by antigen-pulsed
dendritic cells)

L40 ANSWER 12 OF 15 HCAPLUS COPYRIGHT 2004 ACS on STN

AN 1995:842985 HCAPLUS

DN 123:283435

ED Entered STN: 10 Oct 1995

TI Acutely infected Langerhans cells are more efficient than T cells in
disseminating HIV type 1 to **activated** T cells
following a short cell-cell contact

AU Ayehunie, Seyoum; Groves, Richard W.; Bruzzese, Ann-Marie; Ruprecht, Ruth

M.; Kupper, Thomas S.; Langhoff, Erik
 CS Laboratory Viral Pathogenesis, Dana-Farber Cancer Institute, Boston, MA,
 02115, USA
 SO AIDS Research and Human Retroviruses (1995), 11(8), 877-84
 CODEN: ARHRE7; ISSN: 0889-2229
 PB Liebert
 DT Journal
 LA English
 CC 15-8 (Immunochemistry)
 AB Most **human immunodeficiency virus** type 1 (**HIV-1**) infections involve sexual contact and virus passage across mucosal surfaces. While Langerhans cells (LCs) and **dendritic cells** (DCs) have been implicated in mucosal infection, their role is undefined. Here we demonstrate that acutely **HIV-1**-infected LCs and DCs effectively transmit virus to uninfected, **activated** T cells. Cocultivation of these cells results in massive virus production that requires a short cell-cell contact; as little as 30 min contact time is sufficient for **HIV-1-pulsed** DCs to infect their target T cells. Furthermore, surface-bound virus **inactivation** by trypsin does not significantly decrease the efficiency of virus transmission by LC/DCs, suggesting rapid internalization of virus. This effective virus transfer by infected LCs and blood-derived DCs requires prior **activation** of T cells. Surprisingly, cocultivation of acutely infected T cells with uninfected, **activated** target T cells results only in low virus production, even with T cell-tropic virus. We conclude that LCs and DCs are not only important targets of **HIV-1** infection, but may also play a key role in the early dissemination of virus to T cells they encounter in skin or lymphoid tissue.

ST HIV1 virus Langerhans cell **dendritic cell**
 IT Skin, disease
 (Langerhans' cell, infection, role of Langerhan's cells and **dendritic cells** in transmission of **HIV-1** virus to T-cells)
 IT Lymphocyte
 (T-cell, disease, infection, role of Langerhan's cells and **dendritic cells** in transmission of **HIV-1** virus to T-cells)
 IT Leukocyte
 (**dendritic cell**, infection; role of Langerhan's cells and **dendritic cells** in transmission of **HIV-1** virus to T-cells)
 IT Virus, animal
 (**human immunodeficiency 1**, role of Langerhan's cells and **dendritic cells** in transmission of **HIV-1** virus to T-cells)

L40 ANSWER 13 OF 15 HCAPLUS COPYRIGHT 2004 ACS on STN
 AN 1995:308730 HCAPLUS
 DN 122:72000
 ED Entered STN: 24 Jan 1995
 TI Compositions and use of glucocorticoids for treating and preventing AIDS
 IN **Andrieu, Jean-Marie**; Levy, Rafael; Lu, Louis
 PA Association pour la Recherche, l'Etude le Traitement et la Prevention des Maladies Malignes du Sang (AREMAS), Fr.
 SO PCT Int. Appl., 63 pp.
 CODEN: PIXXD2
 DT Patent
 LA French
 IC ICM A61K031-57
 CC 1-5 (Pharmacology)
 Section cross-reference(s): 63
 FAN.CNT 1
 PATENT NO. KIND DATE APPLICATION NO. DATE

 PI WO 9421264 A1 19940929 WO 1994-FR282 19940315
 W: US
 RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE
 FR 2711920 A1 19950512 FR 1993-2966 19930315
 FR 2711920 B1 19960216
 EP 641210 A1 19950308 EP 1994-909969 19940315
 R: BE, CH, DE, ES, FR, GB, IT, LI, NL, SE
 PRAI FR 1993-2966 19930315
 WO 1994-FR282 19940315
 AB At least one glucocorticoid is used to provide a drug for treating HIV infection and preventing AIDS, in particular in a patient infected with HIV-1. A pharmaceutical composition provided for this purpose includes, as the **active** principle, at least one glucocorticoid and/or a salt or other pharmacol. acceptable derivative thereof, particularly prednisone or prednisolone, in a unit dose of approx. 1 mg to 1 g. Results of HIV1-seropos. patients treated with prednisolone are presented. There was a clear increase in the number of CD4 lymphocytes and in the CD4 lymphocyte/CD8 lymphocyte ratio. There was also a reduction in all markers tested for immune **activation** (IgG, IgA, β 2 microglobulin, etc.).
 ST glucocorticoid AIDS treatment; HIV virus infection treatment
 glucocorticoid; prednisolone HIV virus infection treatment
 IT Acquired immune deficiency syndrome
 Immunomodulators
 Pharmaceutical dosage forms
 (compns. and use of glucocorticoids for treating and preventing AIDS)
 IT Antigens
 RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)
 (CD4, CD4 lymphocyte; compns. and use of glucocorticoids for treating and preventing AIDS in relation to CD4 lymphocyte/CD8 lymphocyte ratio)
 IT Antigens
 RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)
 (CD8, CD8 lymphocyte; compns. and use of glucocorticoids for treating and preventing AIDS in relation to CD4 lymphocyte/CD8 lymphocyte ratio)
 IT Corticosteroids, biological studies
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (gluco-, compns. and use of glucocorticoids for treating and preventing AIDS)
 IT Virus, animal
 (human immunodeficiency, compns. and use of glucocorticoids for treating and preventing AIDS)
 IT Virus, animal
 (human immunodeficiency 1, compns. and use of glucocorticoids for treating and preventing AIDS)
 IT 50-02-2, Dexamethasone 50-03-3, Hydrocortisone acetate 50-04-4, Cortisone acetate 50-24-8, Prednisolone 52-21-1, Prednisolone acetate 53-36-1, Methylprednisolone acetate 67-78-7, Triamcinolone diacetate 76-25-5, Triamcinolone acetonide 83-43-2, Methylprednisolone 124-94-7, Triamcinolone 125-04-2 378-44-9, Betamethasone 514-36-3 1177-87-3, Dexamethasone acetate 1499-59-8, Dihydrocortisone acetate 1597-82-6, Paramethasone acetate 2152-44-5, Betamethasone valerate 2375-03-3 2681-16-5 3385-03-3, Flunisolide 5593-20-4, Betamethasone dipropionate 5611-51-8, Triamcinolone hexacetonide 13926-43-7 13926-44-8
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (compns. and use of glucocorticoids for treating and preventing AIDS)

L40 ANSWER 14 OF 15 HCAPLUS COPYRIGHT 2004 ACS on STN
 AN 1993:668923 HCAPLUS
 DN 119:268923
 ED Entered STN: 25 Dec 1993

TI Induction of **CD8+** cytotoxic T lymphocytes by immunization with syngeneic irradiated **HIV-1** envelope derived peptide-**pulsed dendritic cells**
 AU Takahashi, Hidemi; Nakagawa, Yohko; Yokomuro, Kozo; Berzofsky, Jay A.
 CS Dep. Microbiol. Immun., Nippon Med. Sch., Bunkyo-ku, 113, Japan
 SO International Immunology (1993), 5(8), 849-57
 CODEN: INIMEN; ISSN: 0953-8178
 DT Journal
 LA English
 CC 15-10 (Immunochemistry)
 AB Based on the evidence that **CD8+** cytotoxic T cell (CTL) precursors do not appear to distinguish between virus-infected cells and viral peptide-**pulsed** syngeneic cells, the authors have developed methods for priming class I MHC mol. restricted **CD8+** CTL with such peptides without using any adjuvant. The authors were able to prime in vivo such CTL immunity lasting at least 6 mo with a single i.v. injection of syngeneic 2200-3300 rad irradiated class II MHC mol. expressing splenic **dendritic cells** (DC). No foreign serum source was necessary during the **pulsing**. Interestingly, the authors could not generate significant CTL **activity** with unirradiated or low dose (<1100 rad) irradiated spleen cells. Because a) even purified DC required irradiation for optimal **activity**, b) unirradiated B cells did not significantly inhibit the immunization with DC, and c) B cell depletion did not substitute for irradiation, the effect of irradiation might be more to determine homing of the cells than to eliminate interference by B cells. I.v. immunization was much more effective than s.c. or i.p. immunization. CTL generated by this method could kill both peptide-**pulsed** syngeneic targets and targets endogenously expressing the whole gp160 gene. Moreover, **CD8+** CTL could be primed with the minimal 10-residue core peptide (RGPGRAFVTI) for optimal presentation by class I MHC mols. as efficiently as the original p18. Apparently, DC bearing antigenic peptide may prime antigen-specific **CD8+** CTL in vivo.
 ST cytotoxic T lymphocyte **HIV** glycoprotein gp160; **dendritic cell HIV** peptide cytotoxic lymphocyte; **human immunodeficiency virus** peptide cytotoxic lymphocyte
 IT Vaccines
 (for **human immunodeficiency virus**,
 glycoprotein gp160 peptide in relation to)
 IT Lymphocyte
 (T-cell, cytotoxic, priming of, by immunization with **human immunodeficiency virus** envelope peptide-**pulsed dendritic cells**)
 IT Spleen
 (**dendritic cell**, **human immunodeficiency virus** envelope peptide-expressing, immunization with, cytotoxic T-lymphocyte induction by)
 IT Glycoproteins, specific or class
 RL: BIOL (Biological study)
 (gp160env, of **human immunodeficiency virus**, peptide of, **dendritic cells** bearing, immunization with, cytotoxic T-lymphocytes induction by)
 IT **Virus, animal**
 (**human immunodeficiency 1**, glycoprotein gp160 of, peptide of, **dendritic cells** bearing, immunization with, cytotoxic T-lymphocytes induction by)
 IT Radiation
 (ionizing, cytotoxic T-lymphocyte induction by immunization with **human immunodeficiency virus** envelope peptide-**pulsed dendritic cells** in relation to)
 IT 114991-28-5
 RL: BIOL (Biological study)

(of glycoprotein gp160 from **human immunodeficiency virus, dendritic cell** expressing, immunization with, cytotoxic T-lymphocytes induction by)

L40 ANSWER 15 OF 15 HCAPLUS COPYRIGHT 2004 ACS on STN

AN 1992:104176 HCAPLUS

DN 116:104176

ED Entered STN: 20 Mar 1992

TI Primary proliferative and cytotoxic T-cell responses to HIV induced in vitro by human **dendritic cells**

AU Macatonia, S. E.; Patterson, S.; Knight, S. C.

CS Antigen Presentation Res. Group, MRC Clin. Res. Cent., Harrow/Middlesex, HA1 3UJ, UK

SO Immunology (1991), 74(3), 399-406

CODEN: IMMUAM; ISSN: 0019-2805

DT Journal

LA English

CC 15-8 (Immunochemistry)

AB In earlier studies, primary proliferative and cytotoxic T-cell (CTL) responses to influenza virus were produced in vitro by using mouse **dendritic cells** (DC) **pulsed** with virus or viral peptide as the stimulus for syngeneic T cells in 20- μ L hanging-drop cultures. This system was now adapted for producing primary responses with cells from non-immune donors to produce primary proliferative and CTL responses to **human immunodeficiency virus 1 (HIV)** and to HIV peptides in vitro using cells from normal human peripheral blood. All donors in this study were laboratory personnel with no history of HIV infection. DC enriched from peripheral blood were exposed to HIV in vitro and small nos. were added to T lymphocytes. Proliferative responses to virus-infected DC were obtained after 3 days in culture. After 6 days, CTL were obtained that killed virus-infected autologous (but not allogeneic) phytohemagglutinin (PHA)-stimulated blast cells. Proliferative and CTL responses were obtained using cells from random donors expressing a spectrum of major histocompatibility complex (MHC) types but the CTL, once produced, showed killing restricted by the MHC class I type. Treatment of cultures with monoclonal antibody (mAb) to CD4-pos. cells at the beginning of culture blocked the development of both proliferative and CTL responses, but treatment after 5 days had no effect on the CTL **activity**. Treatment with MCA to CD8-pos. cells at the beginning of culture did not block proliferation, but treatment either before or after the 5-day culture period blocked CTL responses. Collaboration between proliferating CD4-pos. cells and CD8-pos. cells may thus be required to produce CTL of the CD8 phenotype. DC exposed to HIV also produced CTL that killed autologous blast cells **pulsed** with gp120 envelope glycoprotein. However, DC infected with whole virus did not produce CTL that lysed target cells **pulsed** with a synthetic peptide, which included a known T-cell epitope of gp120 (representing amino acids 111-126). DC **pulsed** with gp120 were a poor stimulus for the development of CTL. In contrast, DC **pulsed** with the peptide (111-126) stimulated both proliferative and CTL responses. The latter killed not only target cells **pulsed** with the peptide itself or with gp120 but also killed virus-infected autologous blast cells. CTL were again obtained reproducibly with this peptide using donors expressing a spectrum of MHC types. Therefore, cells from donors not infected with HIV and who are not immunocompromised were used to identify a T-cell epitope which, in individuals of different MHC types, initiates the production of CTL which kill virus-infected, target cells. This approach should identify peptides with protective potential for vaccination purposes.

ST HIV peptide cytotoxic lymphocyte **dendritic cell**

IT Lymphocyte
(T-cell, cytotoxic, human immunodeficiency
virus-specific, induction of, by whole virus- vs. viral
peptide-pulsed dendritic cells)

IT Leukocyte
(dendritic cell, human
immunodeficiency virus- or viral peptide-
pulsed, proliferative and cytotoxic T-cell responses induction
by)

IT Sialoglycoproteins
RL: BIOL (Biological study)
(gp120env, peptide of, dendritic cells
pulsed with, of HIV-1 virus, proliferative and
cytotoxic T-cell responses induction by)

IT Virus, animal
(human immunodeficiency 1,
dendritic cells pulsed with viral peptide
vs. whole, proliferative and cytotoxic T-cell responses induction by)

IT 139035-13-5
RL: BIOL (Biological study)
(dendritic cells pulsed with, of
HIV-1 virus, proliferative and cytotoxic T-cell responses
induction by)

=> => fil medline

FILE 'MEDLINE' ENTERED AT 11:19:27 ON 03.FEB 2004

FILE LAST UPDATED: 31 JAN 2004 (20040131/UP). FILE COVERS 1958 TO DATE.

On December 14, 2003, the 2004 MeSH terms were loaded. See HELP RLOAD
for details.

MEDLINE thesauri in the /CN, /CT, and /MN fields incorporate the
MeSH 2004 vocabulary. See <http://www.nlm.nih.gov/mesh/> and
http://www.nih.gov/pubs/yechebull/nd03/nd03_mesh.html for a description
on changes.

This file contains CAS Registry Numbers for easy and accurate
substance identification.

=> d all tot

L87 ANSWER 1 OF 15 MEDLINE on STN
AN 2003440637 MEDLINE
DN 22862988 PubMed ID: 14501788
TI Presentation of exogenous whole inactivated simian
immunodeficiency virus by mature dendritic cells
induces CD4+ and CD8+ T-cell responses.
AU Frank Ines; Santos John J; Mehlhop Erin; Villamide-Herrera Loreley;
Santisteban Christine; Gettie Agegnehu; Ignatius Ralf; Lifson Jeffrey D;
Pope Melissa
CS Center for Biomedical Research, Population Council, New York, New York
10021, USA.
NC AI47681 (NIAID)
AI52060 (NIAID)
N01-CO-12400 (NCI)
RR00164 (NCRR)
SO JOURNAL OF ACQUIRED IMMUNE DEFICIENCY SYNDROMES, (2003 Sep 1) 34 (1) 7-19.
Journal code: 100892005. ISSN: 1525-4135.
CY United States
DT Journal; Article; (JOURNAL ARTICLE)
LA English

FS Priority Journals; AIDS
EM 200310
ED Entered STN: 20030923
Last Updated on STN: 20031010
Entered Medline: 20031009

AB Interactions between **HIV-1** and **dendritic cells**
(DCs) play an important role in the initial establishment and spread of infection and development of antiviral immunity. We used chemically **inactivated** aldrithiol-2 (AT-2) simian immunodeficiency virus (SIV) with functional envelope glycoproteins to study virus interactions with DCs and developed an in vitro system to evaluate the quality of SIV antigen (Ag) presentation by DCs to T cells. AT-2 SIV interacts authentically with T cells and DCs and thus allows assessment of natural SIV-specific responses. CD4+ and CD8+ T cells from blood or lymph nodes of SIV-infected macaques released interferon-gamma (IFN gamma) and proliferated in response to a variety of AT-2 SIV isolates. Responses did not vary significantly as a function of the quantitative envelope glycoprotein content of the virions. Presentation of Ags derived from AT-2 SIV by DCs was more potent than presentation by comparably Ag-loaded monocytes. Interestingly, SIV-pulsed mature DCs stimulated both CD4+ and CD8+ T-cell responses, whereas immature DCs primarily stimulated CD4+ T cells. Further studies using AT-2 **inactivated** virus may help to define better the details of the virus-DC interactions critical for infection versus induction of antiviral immune responses.

CT Check Tags: Animal; Female; Male; Support, Non-U.S. Gov't; Support, U.S. Gov't, P.H.S.
*2,2'-Dipyridyl: AA, analogs & derivatives
2,2'-Dipyridyl: PD, pharmacology
*Antigen Presentation
*CD4-Positive T-Lymphocytes: IM, immunology
*CD8-Positive T-Lymphocytes: IM, immunology
Cell Differentiation
Dendritic Cells: CY, cytology
*Dendritic Cells: IM, immunology
Dendritic Cells: VI, virology
Disulfides: PD, pharmacology
Leukocytes, Mononuclear: IM, immunology
Lymph Nodes: IM, immunology
*Lymphocyte Activation
Macaca mulatta
SAIDS Vaccines: IM, immunology
*SIV: IM, immunology
SIV: PY, pathogenicity
SIV: PH, physiology
Vaccines, Inactivated: IM, immunology

RN 2127-03-9 (2,2'-dipyridyl disulfide); 366-18-7 (2,2'-Dipyridyl).
CN 0 (Disulfides); 0 (SAIDS Vaccines); 0 (Vaccines, **Inactivated**)

L87 ANSWER 2 OF 15 MEDLINE on STN
AN 2003399669 MEDLINE
DN PubMed ID: 12928421
TI Most highly exposed seronegative men lack HIV-1-specific, IFN-gamma-secreting T cells.
AU Hladik Florian; Desbien Anthony; Lang Jean; Wang Lei; Ding Yan; Holte Sarah; Wilson Aaron; Xu Younong; Moerbe Micky; Schmechel Steve; McElrath M Juliana
CS Program in Infectious Diseases, Clinical Research Division, Public Health Sciences Division, Fred Hutchinson Cancer Research Center, 1100 Fairview Avenue North, D3-100, Seattle, WA 98109, USA.
NC AI 27757 (NIAID)
AI 47806 (NIAID)
AI 48017 (NIAID)
SO Journal of immunology (Baltimore, Md. : 1950), (2003 Sep 1) 171 (5)

2671-83.

Journal code: 2985117R. ISSN: 0022-1767.

CY United States

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Abridged Index Medicus Journals; Priority Journals

EM 200311

ED Entered STN: 20030827

Last Updated on STN: 20031218

Entered Medline: 20031117

AB Naturally acquired cellular immunity in individuals who have been exposed to HIV-1 but have remained uninfected may hold clues for the design of an effective HIV vaccine. To determine the presence and nature of such an HIV-1-specific immune response, we evaluated the quantity and fine specificity of HIV-1-reactive IFN-gamma-secreting T cells in a group of highly exposed seronegative men having sex with men. All 46 ES reported frequent unprotected anal sex with known HIV-1-infected partners at enrollment, and high risk activities continued in at least one-half of the volunteers for up to >6 years of observation. Despite the high frequency of unprotected anal intercourse and potential HIV-1 exposure, the vast majority of individuals demonstrated no or very low numbers of HIV-1-specific, IFN-gamma-secreting T cells. Even when HIV-1 epitopes were presented by peptide-pulsed autologous dendritic cells in 15 of the highest risk volunteers, HIV-1-specific T cells remained infrequent, and the proportion of responders was not significantly different from that in a lower risk seronegative control cohort. Only PBMC from two individuals who have remained uninfected to date exhibited distinctly positive responses. However, these responses rarely persisted over time, single epitope specificities were identified in only one volunteer, and HIV-1-specific memory T cell clones did not expand in vitro. HIV-1-specific, IFN-gamma-secreting T cells are thus unlikely to substantially contribute to resistance against infection in most exposed seronegative men having sex with men.

CT Check Tags: Human; Male; Support, Non-U.S. Gov't; Support, U.S. Gov't, P.H.S.

Adolescent

Adult

Amino Acid Sequence

Antigen Presentation

Clone Cells

Dendritic Cells: IM, immunology

Dendritic Cells: ME, metabolism

Enzyme-Linked Immunosorbent Assay: MT, methods

Enzyme-Linked Immunosorbent Assay: ST, standards

Epitope Mapping

Epitopes, T-Lymphocyte: AN, analysis

Epitopes, T-Lymphocyte: IM, immunology

*HIV Infections: IM, immunology

HIV Infections: ME, metabolism

*HIV Seronegativity: IM, immunology

*HIV-1: IM, immunology

*Interferon Type II: SE, secretion

Lymphocyte Count

Middle Aged

Molecular Sequence Data

Risk-Taking

*T-Lymphocyte Subsets: IM, immunology

*T-Lymphocyte Subsets: SE, secretion

T-Lymphocyte Subsets: VI, virology

T-Lymphocytes, Cytotoxic: IM, immunology

T-Lymphocytes, Cytotoxic: SE, secretion

T-Lymphocytes, Cytotoxic: VI, virology

RN 82115-62-6 (Interferon Type II)

CN 0 (Epitopes, T-Lymphocyte)

L87 ANSWER 3 OF 15 MEDLINE on STN

AN 2003364765 MEDLINE

DN 22768196 PubMed ID: 12885891

TI Induction of protective immune responses against R5 **human immunodeficiency virus** type 1 (HIV-1) infection in hu-PBL-SCID mice by intrasplenic immunization with **HIV-1-pulsed dendritic cells**:

possible involvement of a novel factor of human CD4(+) T-cell origin.

AU Yoshida Atsushi; Tanaka Reiko; Murakami Tsutomu; Takahashi Yoshiaki; Koyanagi Yoshio; Nakamura Masataka; Ito Mamoru; Yamamoto Naoki; Tanaka Yuetsu

CS Department of Immunology, Graduate School and Faculty of Medicine, University of the Ryukyus, Nishihara, Okinawa 903-0215, Japan.

SO JOURNAL OF VIROLOGY, (2003 Aug) 77 (16) 8719-28.

Journal code: 0113724. ISSN: 0022-538X.

CY United States

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals

EM 200309

ED Entered STN: 20030806

Last Updated on STN: 20030924

Entered Medline: 20030923

AB The potential of a **dendritic cell** (DC)-based vaccine against **human immunodeficiency virus** type 1 (HIV-1) infection in humans was explored with SCID mice reconstituted with human peripheral blood mononuclear cells (PBMC). HIV-1-negative normal human PBMC were transplanted directly into the spleens of SCID mice (hu-PBL-SCID-spl mice) together with autologous mature DCs **pulsed** with either **inactivated HIV** -1 (strain R5 or X4) or ovalbumin (OVA), followed by a booster injection 5 days later with autologous DCs **pulsed** with the same respective antigens. Five days later, these mice were challenged intraperitoneally with R5 HIV-1(JR-CSF). Analysis of infection at 7 days postinfection showed that the DC-HIV-1-immunized hu-PBL-SCID-spl mice, irrespective of the HIV-1 isolate used for immunization, were protected against HIV-1 infection. In contrast, none of the DC-OVA-immunized mice were protected. Sera from the DC-HIV -1- but not the DC-OVA-immunized mice inhibited the in vitro infection of activated PBMC and macrophages with R5, but not X4, HIV-1. Upon restimulation with HIV-1 in vitro, the human CD4(+) T cells derived from the DC-HIV-1-immunized mice produced a similar R5 HIV-1 suppressor factor. Neutralizing antibodies against human RANTES, MIP-1alpha, MIP-1beta, alpha interferon (IFN-alpha), IFN-beta, IFN-gamma, interleukin-4 (IL-4), IL-10, IL-13, IL-16, MCP-1, MCP-3, tumor necrosis factor alpha (TNF-alpha), or TNF-beta failed to reverse the HIV-1-suppressive activity. These results show that **inactivated HIV-1-pulsed** autologous DCs can stimulate splenic resident human CD4(+) T cells in hu-PBL-SCID-spl mice to produce a yet-to-be-defined, novel soluble factor(s) with protective properties against R5 HIV-1 infection.

CT Check Tags: Animal; Support, Non-U.S. Gov't

AIDS Vaccines: IM, immunology

*CD4-Positive T-Lymphocytes: IM, immunology

Cytokines: IM, immunology

*Dendritic Cells: IM, immunology

*HIV-1: IM, immunology

Mice

Mice, Inbred BALB C

Mice, SCID
Neutralization Tests
*Spleen: IM, immunology
CN 0 (AIDS Vaccines); 0 (Cytokines)

L87 ANSWER 4 OF 15 MEDLINE on STN
AN 2003364691 MEDLINE
DN 22756620 PubMed ID: 12874266
TI Potent immune response against HIV-1 and protection from virus challenge in hu-PBL-SCID mice immunized with **inactivated virus-pulsed dendritic cells** generated in the presence of IFN-alpha.
AU Lapenta Caterina; Santini Stefano M; Logozzi Mariantonia; Spada Massimo; Andreotti Mauro; Di Pucchio Tiziana; Parlato Stefania; Belardelli Filippo
CS Laboratory of Virology, Istituto Superiore di Sanita, Viale Regina Elena, 299, Rome, Italy 00161.
SO JOURNAL OF EXPERIMENTAL MEDICINE, (2003 Jul 21) 198 (2) 361-7.
Journal code: 2985109R. ISSN: 0022-1007.
CY United States
DT Journal; Article; (JOURNAL ARTICLE)
LA English
FS Priority Journals
EM 200309
ED Entered STN: 20030806
Last Updated on STN: 20030926
Entered Medline: 20030925
AB A major challenge of AIDS research is the development of therapeutic vaccine strategies capable of inducing the humoral and cellular arms of the immune responses against HIV-1. In this work, we evaluated the capability of DCs **pulsed** with aldrithiol-2-**inactivated HIV-1** in inducing a protective antiviral human immune response in SCID mice reconstituted with human PBL (hu-PBL-SCID mice). Immunization of hu-PBL-SCID mice with DCs generated after exposure of monocytes to GM-CSF/IFN-alpha (IFN-DCs) and **pulsed with inactivated HIV-1** resulted in a marked induction of human anti-HIV-1 antibodies, which was associated with the detection of anti-HIV neutralizing activity in the serum. This vaccination schedule also promoted the generation of a human CD8+ T cell response against HIV-1, as measured by IFN-gamma Elispot analysis. Notably, when the hu-PBL-SCID mice immunized with antigen-**pulsed** IFN-DCs were infected with HIV-1, inhibition of virus infection was observed as compared with control animals. These results suggest that IFN-DCs **pulsed with inactivated HIV-1** can represent a valuable approach of immune intervention in HIV-1-infected patients.
CT Check Tags: Animal; Human; Support, Non-U.S. Gov't
*AIDS Vaccines: TU, therapeutic use
*Acquired Immunodeficiency Syndrome: IM, immunology
Acquired Immunodeficiency Syndrome: PC, prevention & control
*Dendritic Cells: IM, immunology
Dendritic Cells: TR, transplantation
Dendritic Cells: VI, virology
*HIV-1: IM, immunology
Immunomagnetic Separation: MT, methods
*Interferon-alpha: IM, immunology
Lymphocyte Transfusion
Lymphocytes: CY, cytology
*Lymphocytes: IM, immunology
Mice
Mice, SCID
Transplantation, Heterologous: IM, immunology
*Vaccines, Inactivated: TU, therapeutic use
CN 0 (AIDS Vaccines); 0 (Interferon-alpha); 0 (Vaccines, Inactivated

)

L87 ANSWER 5 OF 15 MEDLINE on STN
AN 2003357445 MEDLINE
DN 22771896 PubMed ID: 12891059
TI **Dendritic cells** generated in the presence of
granulocyte-macrophage colony-stimulating factor and IFN-alpha are potent
inducers of **HIV**-specific CD8 T cells.
AU Carbonneil Cedric; Aouba Albertine; Burgard Marianne; Cardinaud Sylvain;
Rouzioux Christine; Langlade-Demoyen Pierre; Weiss Laurence
CS INSERM U430, Paris, France.
SO AIDS, (2003 Aug 15) 17 (12) 1731-40.
Journal code: 8710219. ISSN: 0269-9370.
CY England: United Kingdom
DT Journal; Article; (JOURNAL ARTICLE)
LA English
FS Priority Journals; AIDS
EM 200310
ED Entered STN: 20030801
Last Updated on STN: 20031008
Entered Medline: 20031007
AB OBJECTIVE: To investigate the ability of granulocyte-macrophage
colony-stimulating factor (GM-CSF) and IFN-alpha to induce the
differentiation of peripheral monocytes into **dendritic**
cells (DC) and their ability to trigger an **HIV**-specific
CD8 T-cell response. METHODS: Monocytes isolated from both seronegative
controls and **HIV**-infected individuals were differentiated into
DC using GM-CSF with either IL-4 or IFN-alpha for 7 days. We assessed the
phenotypic characteristics and IL-12 production by flow cytometry. The
ability of DC to trigger CD8 T-cell responses was assessed by means of
ELISpot and cytotoxicity assays. In addition, **HIV**-1-RNA levels
were measured in culture supernatants. RESULTS: Compared with control DC
generated in the presence of GM-CSF and IL-4, DC generated in the presence
of GM-CSF and IFN-alpha expressed higher levels of MHC class I molecules
and produced similar or higher levels of IL-12 after CD40 ligation or
Staphylococcus aureus Cowan stimulation. GM-CSF/IFN-alpha DC expressed low
levels of CD4, CXCR4 and DC-SIGN and did not produce detectable virus
during the differentiation period. **Pulsed** GM-CSF/IFN-alpha DC
were found to prime CD8 T cells from **HIV**-negative controls to
exert cytotoxic activity against target cells expressing **HIV**
antigens. **HIV** peptide-**pulsed** GM-CSF/IFN-alpha DC
promote specific IFN-gamma production by autologous CD8 T cells from
HIV-seronegative donors. Furthermore, GM-CSF/IFN-alpha DC from
HIV-seropositive patients efficiently present **HIV**
peptides to autologous CD8 T lymphocytes. CONCLUSION: GM-CSF and
IFN-alpha allow the generation of DC with high CD8 T-cell stimulating
abilities. Therefore, this strategy may represent a novel approach to
therapeutic vaccination in **HIV** disease.
CT Check Tags: Comparative Study; Human; Support, Non-U.S. Gov't
*Adoptive Transfer: MT, methods
CD40 Ligand: PD, pharmacology
*CD8-Positive T-Lymphocytes: IM, immunology
Cell Differentiation
Cells, Cultured
*Dendritic Cells: IM, immunology
Granulocyte-Macrophage Colony-Stimulating Factor: IM, immunology
*Granulocyte-Macrophage Colony-Stimulating Factor: PD, pharmacology
HIV Infections: IM, immunology
*HIV Infections: TH, therapy
HIV-1: GE, genetics
HIV-1: IM, immunology
Interferon Type II: IM, immunology
*Interferon Type II: PD, pharmacology

Interleukin-12: IM, immunology
Interleukin-4: PD, pharmacology

Lymphocyte Activation

RNA, Viral: AN, analysis

Staphylococcus aureus

T-Lymphocytes, Cytotoxic: IM, immunology

RN 147205-72-9 (CD40 Ligand); 187348-17-0 (Interleukin-12); 207137-56-2
(Interleukin-4); 82115-62-6 (Interferon Type II); 83869-56-1
(Granulocyte-Macrophage Colony-Stimulating Factor)
CN 0 (RNA, Viral)

L87 ANSWER 6 OF 15 MEDLINE on STN

AN 2003031369 MEDLINE

DN 22426478 PubMed ID: 12538688

TI Induction of antigen-specific CTL by recombinant HIV
trans-activating fusion protein-**pulsed** human monocyte-derived
dendritic cells.

AU Tanaka Yoshiyuki; Dowdy Steven F; Linehan David C; Eberlein Timothy J;
Goedegebuure Peter S

CS Department of Surgery, Biologic Cancer Therapy Program, and Alvin J.
Siteman Cancer Center, Washington University School of Medicine, St.
Louis, MO 63110, USA.

NC K08CA87018 (NCI)

R01CA68500 (NCI)

SO JOURNAL OF IMMUNOLOGY, (2003 Feb 1) 170 (3) 1291-8.

Journal code: 2985117R. ISSN: 0022-1767.

CY United States

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Abridged Index Medicus Journals; Priority Journals

EM 200303

ED Entered STN: 20030123

Last Updated on STN: 20030328

Entered Medline: 20030327

AB Several systems have been tested for introduction of Ags into human
dendritic cells (DC). Most of them to date, however,
are complex and possess limited efficiency. Recent advances in
HIV trans-activating (TAT) fusion protein technology permit
extremely high transduction efficiencies for a majority of mammalian cell
types. Here we report our attempts to develop a simple, but highly
efficient, protocol for loading of antigenic protein into DC using TAT
fusion technology. A TAT-minigene fusion protein was generated, encoding
both the HLA-A2-restricted influenza matrix protein-derived epitope
(GILVFTFTL, Flu-M1) and a melanoma Ag gp100-derived modified epitope
(YLEPGPVTV, G9(280)-9V). In addition, both a TAT-Her2/neu extracellular
domain (ECD) fusion protein and a TAT-green fluorescence protein fusion
protein were generated. Over 95% of DC stained positively for TAT-green
fluorescence protein within 20 min of coculture. DC treated with
TAT-minigene were efficiently recognized by both Flu-M1 and
G9(280)-9V-specific T cells in cytotoxicity assays and IFN-gamma ELISPOT
assays. In contrast, DC **pulsed** with minigene fusion protein
lacking TAT were either poorly recognized or not recognized by the T
cells. DC **pulsed** with TAT-minigene also efficiently induced
Flu-M1-specific T cells from naive lymphocytes. Similarly, DC treated
with TAT-Her2/neu ECD stimulated patient-derived lymphocytes that
specifically recognized Her2/neu(+) ovarian and breast cancer cell lines.
The CTL induced by TAT-Her2/neu ECD-**pulsed** DC specifically
recognized the Her2/neu ECD-derived immunogenic peptide E75 (KIFGSLAFL).
Our data suggest that TAT fusion proteins efficiently transduce DC and
induce Ag-specific T cells. This could prove to be a useful method for
treatment of infectious diseases and cancer.

CT Check Tags: Human; Support, U.S. Gov't, P.H.S.
Cell Line

Cytotoxicity, Immunologic: GE, genetics
 Dendritic Cells: CY, cytology
 *Dendritic Cells: IM, immunology
 Dendritic Cells: ME, metabolism
 Epitopes, T-Lymphocyte: GE, genetics
 *Epitopes, T-Lymphocyte: IM, immunology
 Extracellular Space: GE, genetics
 Extracellular Space: PH, physiology
 *Gene Products, tat: GE, genetics
 Gene Products, tat: PH, physiology
 *HIV-1: GE, genetics
 HIV-1: IM, immunology
 HLA-A2 Antigen: IM, immunology
 K562 Cells
 *Lymphocyte Activation
 Lymphocyte Activation: GE, genetics
 Membrane Glycoproteins: GE, genetics
 Membrane Glycoproteins: PH, physiology
 Monocytes: CY, cytology
 *Monocytes: IM, immunology
 Neoplasm Proteins: GE, genetics
 Neoplasm Proteins: PH, physiology
 Protein Denaturation
 Protein Structure, Tertiary: GE, genetics
 Receptor, erbB-2: GE, genetics
 Receptor, erbB-2: PH, physiology
 Recombinant Fusion Proteins: IP, isolation & purification
 Recombinant Fusion Proteins: ME, metabolism
 *Recombinant Fusion Proteins: PH, physiology
 *T-Lymphocytes, Cytotoxic: IM, immunology
 Transduction, Genetic
 Tumor Cells, Cultured
 Viral Matrix Proteins: GE, genetics
 Viral Matrix Proteins: PH, physiology
 CN 0 (Epitopes, T-Lymphocyte); 0 (Gene Products, tat); 0 (HLA-A2 Antigen); 0
 (Membrane Glycoproteins); 0 (Neoplasm Proteins); 0 (Recombinant Fusion
 Proteins); 0 (Viral Matrix Proteins); 0 (influenza virus membrane
 protein); 0 (melanocyte lineage-specific antigen gp100); EC 2.7.1.112
 (Receptor, erbB-2)

L87 ANSWER 7 OF 15 MEDLINE on STN
 AN 2003008183 MEDLINE
 DN 22402162 PubMed ID: 12496959
 TI Therapeutic **dendritic-cell** vaccine for simian AIDS.
 CM Comment in: Nat Med. 2003 Jan;9(1):13-4
 AU Lu Wei; Wu Xiaoxian; Lu Yaozeng; Guo Weizhong; **Andrieu Jean-Marie**
 CS Institut de Recherche sur les Vaccins et l'Immunotherapie des Cancers et
 du Sida, Paris, France.. louis.wei-lu@biomedicale.univ-paris5.fr
 SO NATURE MEDICINE, (2003 Jan) 9 (1) 27-32.
 Journal code: 9502015. ISSN: 1078-8956.
 CY United States
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Priority Journals
 EM 200303
 ED Entered STN: 20030107
 Last Updated on STN: 20030319
 Entered Medline: 20030318
 AB An effective immune response against **human**
immunodeficiency virus or simian immunodeficiency virus
 (SIV) is critical in achieving control of viral replication. Here, we
 show in SIV-infected rhesus monkeys that an effective and durable
 SIV-specific cellular and humoral immunity is elicited by a vaccination

with chemically **inactivated SIV-pulsed dendritic cells**. After three immunizations made at two-week intervals, the animals exhibited a 50-fold decrease of SIV DNA and a 1,000-fold decrease of SIV RNA in peripheral blood. Such reduced viral load levels were maintained over the remaining 34 weeks of the study. Molecular and cellular analyses of axillary and inguinal node lymphocytes of vaccinated monkeys revealed a correlation between decreased SIV DNA and RNA levels and increased SIV-specific T-cell responses. Neutralizing antibody responses were augmented and remained elevated.

Inactivated whole virus-pulsed dendritic cell vaccines are promising means to control diseases caused by immuno- deficiency viruses.

- CT Check Tags: Animal; Female; Human; Male; Support, Non-U.S. Gov't
 *2,2'-Dipyridyl: AA, analogs & derivatives
 2,2'-Dipyridyl: ME, metabolism
 AIDS Vaccines: IM, immunology
 AIDS Vaccines: TU, therapeutic use
 Antibodies, Viral: BL, blood
 CD4 Lymphocyte Count
 CD4-Positive T-Lymphocytes: IM, immunology
 Dendritic Cells: CY, cytology
 *Dendritic Cells: IM, immunology
 Dendritic Cells: ME, metabolism
 Disulfides: ME, metabolism
 HIV Infections: IM, immunology
 HIV Infections: TH, therapy
 Immunity, Cellular
 Lymph Nodes: IM, immunology
 Lymph Nodes: PA, pathology
 Lymph Nodes: VI, virology
 Macaca mulatta
 Neutralization Tests
 Oxidants: ME, metabolism
 RNA, Viral: AN, analysis
 SAIDS Vaccines: IM, immunology
 *SAIDS Vaccines: TU, therapeutic use
 *SIV: IM, immunology
 SIV: PH, physiology
 Simian Acquired Immunodeficiency Syndrome: IM, immunology
 *Simian Acquired Immunodeficiency Syndrome: TH, therapy
 T-Lymphocytes, Cytotoxic: IM, immunology
 Vaccination
 Viral Load
- RN 2127-03-9 (2,2'-dipyridyl disulfide); 366-18-7 (2,2'-Dipyridyl)
- CN 0 (AIDS Vaccines); 0 (Antibodies, Viral); 0 (Disulfides); 0 (Oxidants); 0 (RNA, Viral); 0 (SAIDS Vaccines)
- L87 ANSWER 8 OF 15 MEDLINE on STN
- AN 2002394649 MEDLINE
- DN 22121899 PubMed ID: 12131208
- TI Activation of HIV-1 specific CD4 and CD8 T cells by human **dendritic cells**: roles for cross-presentation and non-infectious HIV-1 virus.
- AU Larsson Marie; Fonteneau Jean-Francois; Lirvall Margareta; Haslett Patrick; Lifson Jeffrey D; Bhardwaj Nina
- CS The Laboratory of Cellular Physiology and Immunology, The Rockefeller University, New York 10021, USA.
- NC AI 44628 (NIAID)
 AI 47742 (NIAID)
 M01-RR00102 (NCRR)
 N01-CO-56000 (NCI)
- SO AIDS, (2002 Jul 5) 16 (10) 1319-29.
 Journal code: 8710219. ISSN: 0269-9370.

CY England: United Kingdom
DT Journal; Article; (JOURNAL ARTICLE)
LA English
FS Priority Journals; AIDS
EM 200302
ED Entered STN: 20020730
Last Updated on STN: 20030226
Entered Medline: 20030225

AB BACKGROUND: The CD4 T cells in mucosal subepithelia are the first cells to become infected during sexual transmission of HIV-1. **Dendritic cells** (DC) are located in the same area and are known to play a central role in antiviral immune responses. However, extensive viral replication, syncytia formation and cell death follows the interaction between T cells and DC previously exposed to HIV-1. Despite this, anti-HIV responses are generated that control viremia following acute infection. OBJECTIVE: The anti-HIV-1 cellular immune responses observed may be activated by sources other than productively infected DC. HIV-1 induces apoptosis both in cells it infects and in bystander cells. Furthermore, retroviral replication typically generates a predominance of defective particles. We tested whether DC exposed to antigen from either of these sources could elicit anti-HIV specific immune responses. DESIGN AND METHODS: Apoptotic or necrotic monocytes infected with vaccinia virus vectors encoding HIV antigens, a cell line with integrated HIV -1 and apoptotic CD4 T cells **pulsed** with non-infectious or infectious HIV-1 virus were used as sources of antigens to assess cross presentation by DC. Furthermore, direct DC presentation of antigen from non-infectious and infectious HIV-1 was examined. RESULTS: We find that dead cells expressing HIV-1 antigens as well as non-infectious HIV-1 particles can be acquired and processed by DC, leading to the activation, differentiation and expansion of viral antigen-specific CD4 and CD8 T cells from seropositive individuals. CONCLUSIONS: These sources of antigens may be critical for the generation and maintenance of anti-HIV-1 immunity by DC.

CT Check Tags: Human; Support, Non-U.S. Gov't; Support, U.S. Gov't, P.H.S.
*Antigen Presentation
Apoptosis
*CD4-Positive T-Lymphocytes: IM, immunology
CD4-Positive T-Lymphocytes: VI, virology
*CD8-Positive T-Lymphocytes: IM, immunology
CD8-Positive T-Lymphocytes: VI, virology
*Dendritic Cells: IM, immunology
Dendritic Cells: PA, pathology
HIV Antigens: IM, immunology
*HIV-1: IM, immunology
Hela Cells
Immunity, Cellular
Immunologic Memory
Lymphocyte Activation

CN 0 (HIV Antigens)

L87 ANSWER 9 OF 15 MEDLINE on STN
AN 2001694051 MEDLINE
DN 21605992 PubMed ID: 11738745
TI Protection against chronic infection and AIDS by an HIV envelope peptide-cocktail vaccine in a pathogenic SHIV-rhesus model.
AU Nehete P N; Chitta S; Hossain M M; Hill L; Bernacky B J; Baze W; Arlinghaus R B; Sastry K J
CS Department of Veterinary Sciences, The University of Texas M.D. Anderson Cancer Center, Science Park, 650 Coolwater Drive, Bastrop, TX 78602, USA.
NC AI 42694 (NIAID)
CA 16672 (NCI)
SO VACCINE, (2001 Dec 12) 20 (5-6) 813-25.

Journal code: 8406899. ISSN: 0264-410X.

CY England: United Kingdom

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals

EM 200204

ED Entered STN: 20011217

Last Updated on STN: 20020413

Entered Medline: 20020412

AB Based on our prior studies in mouse, monkey, chimpanzee, and human experimental systems, we identified six peptides encoded by highly conserved regions of the **human immunodeficiency virus** type 1 (HIV-1) envelope gene that selectively induce cellular immune responses in the absence of anti-viral antibody production. We tested a cocktail of the six peptides as a prototype vaccine for protection from simian **human immunodeficiency virus** (SHIV) infection and acquired immunodeficiency syndrome (AIDS) in a rhesus monkey model. Three monkeys were vaccinated with the peptide cocktail in Freund's adjuvant followed by autologous **dendritic cells** (DC) **pulsed** with these peptides. All the vaccinated animals exhibited significant induction of T-cell proliferation and cytotoxic T lymphocytes (CTL) responses, but no neutralizing antibodies. Two control mock-vaccinated monkeys showed no specific immune responses. Upon challenge with the pathogenic SHIV(KU-2), both the control and vaccinated monkeys were infected, but efficient clearance of virus-infected cells was observed in all the three vaccinated animals within 14 weeks. These animals also experienced a boosting of antiviral cellular immune responses after infection, and maintained antigen-specific IFN-gamma-producing cells in circulation beyond 42 weeks post-challenge. In contrast, the two mock-vaccinated monkeys had low to undetectable cellular immune responses and maintained significant levels of viral-infected cells and infectious virus in circulation. Further, in both the control monkeys plasma viremia was detectable beyond 38 weeks post-challenge indicating chronic phase infection. In one control monkey, the CD4+ cells dropped to very low levels by 2 weeks post-challenge and became undetectable by week 39 coinciding with high plasma viremia and AIDS, which included cachexia and ataxia. These results serve as proof of principle for the effectiveness of the HIV envelope peptide cocktail vaccine against chronic infection and AIDS, and support the development of multivalent peptide-based vaccine as a viable strategy to induce cell-mediated immunity (CMI) for protection against HIV and AIDS in humans.

CT Check Tags: Animal; Female; Human; Support, U.S. Gov't, P.H.S.

AIDS Vaccines: GE, genetics

AIDS Vaccines: IM, immunology

*AIDS Vaccines: PD, pharmacology

Acquired Immunodeficiency Syndrome: IM, immunology

Acquired Immunodeficiency Syndrome: PC, prevention & control

Amino Acid Sequence

CD4 Lymphocyte Count

HIV Antibodies: BI, biosynthesis

HIV Infections: IM, immunology

HIV Infections: PC, prevention & control

HIV-1: GE, genetics

*HIV-1: IM, immunology

HIV-1: PY, pathogenicity

Immunity, Cellular

Interferon Type II: BI, biosynthesis

Lymphocyte Activation

Macaca mulatta

Mice

SAIDS Vaccines: GE, genetics

SAIDS Vaccines: IM, immunology

*SAIDS Vaccines: PD, pharmacology
 SIV: GE, genetics
 *SIV: IM, immunology
 SIV: PY, pathogenicity
 Simian Acquired Immunodeficiency Syndrome: IM, immunology
 Simian Acquired Immunodeficiency Syndrome: PC, prevention & control
 T-Lymphocytes: IM, immunology
T-Lymphocytes, Cytotoxic: IM, immunology
 Vaccines, Synthetic: GE, genetics
 Vaccines, Synthetic: IM, immunology
 Vaccines, Synthetic: PD, pharmacology
 Viral Envelope Proteins: GE, genetics
 Viral Envelope Proteins: IM, immunology
 RN 82115-62-6 (Interferon Type II)
 CN 0 (AIDS Vaccines); 0 (**HIV** Antibodies); 0 (SAIDS Vaccines); 0
 (Vaccines, Synthetic); 0 (Viral Envelope Proteins)

L87 ANSWER 10 OF 15 MEDLINE on STN
 AN 2001490768 MEDLINE
 DN 21424479 PubMed ID: 11533158
 TI In vitro **human immunodeficiency virus**
 eradication by autologous CD8(+) T cells expanded with **inactivated**
-virus-pulsed dendritic cells.
 AU Lu W; Andrieu J M
 CS Laboratory of Molecular Oncology and Virology, Necker Faculty of Medicine
 at Saints-Peres Biomedical Center, Rene Descartes University, Paris,
 France.. louis.weilu@biomedicale.univ-paris5.fr
 SO JOURNAL OF VIROLOGY, (2001 Oct) 75 (19) 8949-56.
 Journal code: 0113724. ISSN: 0022-538X.
 CY United States
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Priority Journals
 EM 200110
 ED Entered STN: 20010905
 Last Updated on STN: 20011015
 Entered Medline: 20011011
 AB Despite significant immune recovery with potent highly active
 antiretroviral therapy (HAART), eradication of **human**
immunodeficiency virus (HIV) from the bodies
 of infected individuals represents a challenge. We hypothesized that an
 inadequate or inappropriate signal in virus-specific antigen presentation
 might contribute to the persistent failure to mount efficient anti-
HIV immunity in most **HIV**-infected individuals. Here, we
 conducted an in vitro study with untreated (n = 10) and HAART-treated (n =
 20) **HIV** type 1 (**HIV**-1) patients which showed that
pulsing of monocyte-derived dendritic cells
 (DC) with aldrithiol-2-**inactivated** autologous virus resulted in
 the expansion of virus-specific CD8(+) T cells which were capable of
 killing **HIV**-1-infected cells and eradicating the virus from
 cultured patient peripheral blood mononuclear cells independently of the
 disease stages and HAART response statuses of the patients. This in vitro
 anti-**HIV** effect was further enhanced by the **HIV**
 protease inhibitor indinavir (at a nonantiviral concentration), which has
 been shown previously to be able to up-regulate directly patient T-cell
 proliferation following immune stimulation. However, following a 2-day
 treatment with culture supernatant derived from immune-activated T cells
 (which mimics an in vivo environment of **HIV**-disseminated and
 immune-activated lymphoid tissues), DC lost their capacity to present de
 novo **inactivated**-virus-derived antigens. These findings provide
 important information for understanding the establishment of chronic
HIV infection and indicate a perspective for clinical use of
 DC-based therapeutic vaccines against **HIV**.

CT Check Tags: Female; Human; Male; Support, Non-U.S. Gov't
Anti-HIV Agents: IM, immunology
Anti-HIV Agents: TU, therapeutic use
*CD8-Positive T-Lymphocytes: IM, immunology
CD8-Positive T-Lymphocytes: PA, pathology
*Dendritic Cells: IM, immunology
Dendritic Cells: PA, pathology
HIV Infections: DT, drug therapy
*HIV Infections: IM, immunology
HIV Infections: PA, pathology
Immunity, Cellular

CN 0 (Anti-HIV Agents)

L87 ANSWER 11 OF 15 MEDLINE on STN
AN 2001464747 MEDLINE
DN 21400872 PubMed ID: 11509641
TI Enhanced **dendritic cell**-driven proliferation and anti-
HIV activity of CD8(+) T cells by a new phenothiazine derivative,
aminoperazine.
AU Lu W; Achour A; Arlie M; Cao L; **Andrieu J M**
CS Laboratory of Molecular Oncology and Virology, Necker Faculty of Medicine,
Saints-Peres Biomedical Center, Rene Descartes University, Paris, France..
louis.weilu@biomedicale.univ.paris5.fr
SO JOURNAL OF IMMUNOLOGY, (2001 Sep 1) 167 (5) 2929-35.
Journal code: 2985117R. ISSN: 0022-1767.
CY United States
DT Journal; Article; (JOURNAL ARTICLE)
LA English
FS Abridged Index Medicus Journals; Priority Journals
EM 200112
ED Entered STN: 20010820
Last Updated on STN: 20020122
Entered Medline: 20011205

AB T cell anergy, apoptosis, and chronic activation of T lymphocytes are
prevailing features of HIV infection. The inability to develop
an efficient natural antiviral activity in infected patients might be the
consequence of a failure of the Ag presentation by **dendritic**
cells (DCs) in chronically activated lymphoid tissues. We have
identified a new phenothiazine derivative aminoperazine (APR;
2-amino-10-[3'-(1-methyl-4-piperazinyl)propyl]phenothiazine,
C(20)H(26)N(4)S; m.w. 354.51) able to increase (effective dose from 0.1 to
100 nM) the Ag-specific DC-driven proliferation and differentiation of in
vitro HIV-infected and uninfected normal donor T cells and of T
cells from HIV-1-infected patients. The immunomodulatory effect
of APR-sensitized DCs were ascribed to soluble factors derived from DCs.
APR was also capable of increasing HIV gag-p24-specific
proliferation and anti-HIV cytotoxic activity of patients'
CD8(+) T cells against autologous B-lymphoblastoid cell lines expressing a
HIV gag gene, resulting in the suppression of both proviral DNA
and supernatant viral RNA in the HIV-1-infected patients' T cell
culture. This new phenothiazine derivative (APR) might be used for
boosting the immune response of vaccinated individuals and for restoring
the immunity of immunocompromised patients.

CT Check Tags: Human; In Vitro; Support, Non-U.S. Gov't
Adjuvants, Immunologic: CH, chemistry
*Adjuvants, Immunologic: PD, pharmacology
Apoptosis
CD8-Positive T-Lymphocytes: CY, cytology
*CD8-Positive T-Lymphocytes: DE, drug effects
*CD8-Positive T-Lymphocytes: IM, immunology
Cell Division: DE, drug effects
Cell Line
Cytotoxicity, Immunologic: DE, drug effects

*Dendritic Cells: DE, drug effects

*Dendritic Cells: IM, immunology

Genes, gag

HIV Core Protein p24: AD, administration & dosage

HIV Core Protein p24: IM, immunology

HIV Infections: DT, drug therapy

HIV Infections: IM, immunology

HIV-1: GE, genetics

*HIV-1: IM, immunology

Lymphocyte Activation: DE, drug effects

Phenothiazines

Piperazines

Tetradecanoylphorbol Acetate: PD, pharmacology

RN 16561-29-8 (Tetradecanoylphorbol Acetate)

CN 0 (2-amino-10-(3'-(1-methyl-4-piperazinyl)propyl)phenothiazine); 0
(Adjuvants, Immunologic); 0 (HIV Core Protein p24); 0
(Phenothiazines); 0 (Piperazines)

L87 ANSWER 12 OF 15 MEDLINE on STN

AN 1999244883 MEDLINE

DN 99244883 PubMed ID: 10227993

TI Induction of primary human CD8+ T lymphocyte responses in vitro using
dendritic cells.

AU Zarling A L; Johnson J G; Hoffman R W; Lee D R

CS Department of Molecular Microbiology and Immunology, University of
Missouri School of Medicine, Columbia 65212, USA.

NC 5T32 GM08396 (NIGMS)

SO JOURNAL OF IMMUNOLOGY, (1999 May 1) 162 (9) 5197-204.

Journal code: 2985117R. ISSN: 0022-1767.

CY United States

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Abridged Index Medicus Journals; Priority Journals; AIDS

EM 199905

ED Entered STN: 19990601

Last Updated on STN: 19990601

Entered Medline: 19990520

AB The ability of two different human professional APCs, specifically
macrophages (Mphi) and **dendritic cells** (DC), to
stimulate primary responses in human CD8+ T lymphocytes was examined using
both allogeneic and Ag-pulsed autologous APCs. CTL responses in
CD8+ T lymphocytes isolated from HIV-uninfected donors were
evaluated against six different HIV epitopes that are restricted
by four different HLA alleles using autologous human PBMC-derived Mphi and
DCs for primary stimulation. In a side-by-side experiment, immature DCs,
but not Mphi, were able to prime a CTL response against the B14-restricted
p24gag 298-306 epitope; mature DCs were also able to prime a response
against this epitope. In addition, DCs were capable of priming CD8+ CTL
responses against the B8-restricted p24gag 259-267 epitope. In contrast,
Mphi were unable to prime strong CTL responses against other epitopes.
Since the Ag-specific cytotoxic responses required subsequent rounds of
restimulation before they could be detected, the ability of the allogeneic
Mphi and DCs to directly prime CD8+ T lymphocyte responses without
subsequent restimulation was examined. Similar to the aforementioned
peptide-specific results, DCs were more efficient than Mphi in priming
both allogeneic proliferative and cytotoxic responses in human CD8+ T
lymphocytes. Collectively, these results promote an enhanced status for
DCs in the primary stimulation of human CD8+ T lymphocytes.

CT Check Tags: Comparative Study; Human; Support, Non-U.S. Gov't; Support,
U.S. Gov't, P.H.S.

*CD8-Positive T-Lymphocytes: IM, immunology

CD8-Positive T-Lymphocytes: ME, metabolism

Cell Line, Transformed

Cell Separation

Cytotoxicity Tests, Immunologic

***Dendritic Cells: IM, immunology**

Epitopes, T-Lymphocyte: IM, immunology

Epitopes, T-Lymphocyte: ME, metabolism

HIV: IM, immunology

HLA Antigens: GE, genetics

HLA Antigens: ME, metabolism

Histocompatibility Antigens Class I: GE, genetics

Histocompatibility Antigens Class I: ME, metabolism

Isoantigens: GE, genetics

Isoantigens: IM, immunology

***Lymphocyte Activation**

Macrophages: IM, immunology

Protein Binding: IM, immunology

T-Lymphocytes, Cytotoxic: IM, immunology**T-Lymphocytes, Cytotoxic: ME, metabolism****T-Lymphocytes, Cytotoxic: VI, virology**

CN 0 (Epitopes, T-Lymphocyte); 0 (HLA Antigens); 0 (Histocompatibility Antigens Class I); 0 (Isoantigens)

L87 ANSWER 13 OF 15 MEDLINE on STN

AN 1998031752 MEDLINE

DN 98031752 PubMed ID: 9366424

TI Cultured blood **dendritic cells** retain **HIV-1**

antigen-presenting capacity for memory CTL during progressive **HIV-1** infection.

AU Fan Z; Huang X L; Zheng L; Wilson C; Borowski L; Liebmann J; Gupta P; Margolick J; Rinaldo C

CS University of Pittsburgh Graduate School of Public Health, PA 15261, USA.

NC U01-AI-35041 (NIAID)

U01-AI-35042 (NIAID)

U01-AI-37984 (NIAID)

+

SO JOURNAL OF IMMUNOLOGY, (1997 Nov 15) 159 (10) 4973-82.

Journal code: 2985117R. ISSN: 0022-1767.

CY United States

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Abridged Index Medicus Journals; Priority Journals; AIDS

EM 199711

ED Entered STN: 19971224

Last Updated on STN: 19971224

Entered Medline: 19971125

AB **Dendritic cells** (DC) are potent APC that may be involved in the pathogenesis of **HIV-1** infection. We studied the APC function of DC from **HIV-1**-infected subjects that were derived from monocyte-depleted PBMC by culture in human IL-4 and human granulocyte-macrophage CSF. The cultured cells from the **HIV-1**-infected subjects had similar morphology and phenotype of mature DC (CD80 = 41 +/- 8%, CD86 = 77 +/- 5%, CD40 = 87 +/- 6%, CD1a = 1 +/- 1%) to DC cultured from seronegative subjects. The yield of these DC was lower than from **HIV-1**-seronegative subjects (4 +/- 0% vs 11 +/- 2%, p < 0.01), and the lower DC yields correlated with lower numbers of blood CD4+ T cells (r = 0.60, p < 0.01) and higher plasma viral load (r = -0.49, p < 0.01). DC from **HIV-1**-infected subjects were infected with recombinant vaccinia virus vectors expressing Gag, Pol, and Env and were able to stimulate equal or higher levels of MHC class I-restricted, anti-**HIV-1** memory CTL (CTLm) than were similarly treated, autologous B lymphocyte cell lines. DC **pulsed** with peptides representing **HIV-1** CTL epitopes stimulated higher levels of anti-**HIV-1** CTLm responses than did DC infected with the vaccinia virus-**HIV-1** constructs. Allogeneic, MHC class I-matched DC also stimulated anti-

HIV-1 CTLm activity in cells from HIV-1-infected subjects. DC from early and late stages of HIV-1 infection had a similar ability to activate CTLm specific for targets expressing either HIV-1 genes via vaccinia virus vectors or HIV-1 immunodominant synthetic peptides. However, DC from either early or late stages of HIV-1 infection could not overcome the defect in anti-HIV-1 CTLm response in advanced infection.

CT Check Tags: Human; Male; Support, Non-U.S. Gov't; Support, U.S. Gov't, P.H.S.

Acquired Immunodeficiency Syndrome: BL, blood

*Acquired Immunodeficiency Syndrome: IM, immunology

Adult

*Antigen Presentation

Cell Separation: MT, methods

Cells, Cultured

Cytotoxicity, Immunologic: DE, drug effects

*Dendritic Cells: IM, immunology

Dendritic Cells: ME, metabolism

Disease Progression

*HIV Antigens: BL, blood

HIV Antigens: IM, immunology

HIV Seronegativity

HIV Seropositivity: BL, blood

*HIV-1: IM, immunology

Histocompatibility Antigen Class I: AN, analysis

Histocompatibility Testing

*Immunologic Memory

Immunologic Memory: DE, drug effects

Lymphocyte Activation

Middle Age

Oligopeptides: IM, immunology

Oligopeptides: PD, pharmacology

T-Lymphocytes, Cytotoxic: DE, drug effects

*T-Lymphocytes, Cytotoxic: IM, immunology

CN 0 (HIV Antigens); 0 (Histocompatibility Antigen Class I); 0 (Oligopeptides)

L87 ANSWER 14 OF 15 MEDLINE on STN

AN 97265565 MEDLINE

DN 97265565 PubMed ID: 9111471

TI Primary proliferative responses to peptides of HIV Gag p24.

AU Bedford P A; Clarke L B; Hastings G Z; Knight S C

CS Antigen Presentation Research Group, Imperial College School of Medicine, Northwick Park Institute for Medical Research, Harrow, England, U.K.

SO JOURNAL OF ACQUIRED IMMUNE DEFICIENCY SYNDROMES AND HUMAN RETROVIROLOGY, (1997 Apr 1) 14 (4) 301-6.

Journal code: 9501482. ISSN: 1077-9450.

CY United States

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals; AIDS

EM 199705

ED Entered STN: 19970514

Last Updated on STN: 19970514

Entered Medline: 19970506

AB Primary proliferative responses can be initiated by adding

dendritic cells pulsed with antigen to

autologous T cells in 20-microliter hanging drop cultures. To identify primary T-cell epitopes of HIV gag, a series of 23 overlapping peptides, 15 amino acids long, spanning the p24 region were used.

Significant proliferative responses were induced in cells from healthy HIV-negative donors by 11 of these peptides. One of two peptides that bound human leukocyte antigen (HLA)-A *0201 in a peptide-binding

assay using the antigen-processing defective cell line T2 also induced a primary proliferative response. Primary T-cell proliferation was seen in response to some peptides of gag that have not previously been identified as T-cell epitopes in cells from infected individuals. These epitopes might be useful not only for vaccines in antigenically naive individuals but also might increase the breadth of immune responses in seropositive patients.

CT Check Tags: Human; In Vitro

Amino Acid Sequence

HIV Core Protein p24: CH, chemistry

*HIV Core Protein p24: IM, immunology

HIV Seronegativity

*Lymphocyte Activation

Molecular Sequence Data

*Peptides: IM, immunology

*T-Lymphocytes, Cytotoxic: IM, immunology

CN 0 (HIV Core Protein p24); 0 (Peptides)

L87 ANSWER 15 OF 15 MEDLINE on STN

AN 94300111 MEDLINE

DN 94300111 PubMed ID: 8027569

TI Generation of antigen-specific CD8+ CTLs from naive precursors.

AU Mehta-Damani A; Markowicz S; Engleman E G

CS Department of Pathology, Stanford University School of Medicine, CA 94305.

NC AI34313 (NIAID)

HL33811 (NHLBI)

SO JOURNAL OF IMMUNOLOGY, (1994 Aug 1) 153 (3) 996-1003.

Journal code: 2985117R. ISSN: 0022-1767.

CY United States

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Abridged Index Medicus Journals; Priority Journals; AIDS

EM 199408

ED Entered STN: 19940818

Last Updated on STN: 19970203

Entered Medline: 19940809

AB Class I MHC-restricted CTLs are an important component of the host immune response against viral infections, and CTL effectors can often be isolated from infected individuals. However, the mechanism responsible for the induction of CTLs is incompletely understood because, in part, of the difficulty in generating such cells in vitro from naive precursors. In the present study we have used human peripheral blood **dendritic cells** (DCs), devoid of CD4+ T cells, to sensitize naive CD8+ T cells to exogenous Ags, resulting in the generation of Ag specific CTL effectors. With this system, Ag-specific CTL lines were generated to a complex glycoprotein, keyhole limpet hemocyanin, and to multiple small (9-15 amino acids) synthetic peptides derived from conserved regions of the HIV-1 gag and envelope proteins. The HIV-1-specific CTLs demonstrated potent HLA class I restricted killing of both Ag **pulsed** and virally infected target cells. In contrast to Ag-**pulsed** DCs, Ag-**pulsed** monocytes failed to sensitize CTL precursors although they could be used as feeders for purposes of CTL expansion and as target cells in cytolytic assays. With the use of the system described herein, a detailed analysis of the primary human T cell response to foreign Ags is now feasible, and CTL of desired specificity can be generated for potential clinical use in adoptive immunotherapy protocols.

CT Check Tags: Human; In Vitro; Support, U.S. Gov't, Non-P.H.S.; Support, U.S. Gov't, P.H.S.

Antigen-Presenting Cells: IM, immunology

Antigens, CD8: AN, analysis

Cell Separation

Cells, Cultured

Cytotoxicity, Immunologic
 Dendritic Cells: IM, immunology
 HIV Antigens: IM, immunology
 HIV-1: IM, immunology
 Immunity, Cellular

Macrophages: IM, immunology
 Peptides: IM, immunology
 *T-Lymphocyte Subsets: IM, immunology

*T-Lymphocytes, Cytotoxic: IM, immunology

CN 0 (Antigens, CD8); 0 (HIV Antigens); 0 (Peptides)

=> => => d all 27 28 29 35 36

L98 ANSWER 27 OF 47 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN

AN 2002:437885 BIOSIS

DN PREV200200437885

TI Differential transmission of **human immunodeficiency virus** type 1 by distinct subsets of effector **dendritic cells**.

AU Sanders, Rogier W.; de Jong, Esther C.; Baldwin, Christopher E.; Schuitemaker, Joost H. N.; Kapsenberg, Martien L.; Berkhout, Ben [Reprint author]

CS Department of Human Retrovirology, Academic Medical Center, University of Amsterdam, Meibergdreef 15, 1105 AZ, Amsterdam, Netherlands
 b.berkhout@amc.uva.nl

SO Journal of Virology, (August, 2002) Vol. 76, No. 15, pp. 7812-7821. print.
 CODEN: JOVIAM. ISSN: 0022-538X.

DT Article

LA English

ED Entered STN: 14 Aug 2002

Last Updated on STN: 14 Aug 2002

AB **Dendritic cells** (DC) support **human**

immunodeficiency virus type 1 (HIV-1)

transmission by capture of the virus particle in the mucosa and subsequent transport to the draining lymph node, where **HIV-1** is presented to CD4+ Th cells. Virus transmission involves a high-affinity interaction between the DC-specific surface molecule DC-SIGN and the viral envelope glycoprotein gp120 and subsequent internalization of the virus, which remains infectious. The mechanism of viral transmission from DC to T cells is currently unknown. Sentinel immature DC (iDC) develop into Th1-promoting effector DC1 or Th2-promoting DC2, depending on the activation signals. We studied the ability of these effector DC subsets to support **HIV-1** transmission in vitro. Compared with iDC, virus transmission is greatly upregulated for the DC1 subset, whereas DC2 cells are **inactive**. Increased transmission by DC1 correlates with increased expression of ICAM-1, and blocking studies confirm that ICAM-1 expression on DC is important for **HIV** transmission. The ICAM-1-LFA-1 interaction is known to be important for immunological cross talk between DC and T cells, and our results indicate that this cell-cell contact is exploited by **HIV-1** for efficient transmission.

CC Cytology - Animal 02506

Biochemistry studies - General 10060

Blood - Blood and lymph studies 15002

Blood - Blood cell studies 15004

Virology - Animal host viruses 33506

Immunology - General and methods 34502

Medical and clinical microbiology - Virology 36006

IT Major Concepts

Biochemistry and Molecular Biophysics; Immune System (Chemical Coordination and Homeostasis); Infection

IT Parts, Structures, & Systems of Organisms

CD4 positive T cells: immune system; **dendritic cells**

: immune system; lymph node: blood and lymphatics, immune system

ORGN Classifier

Retroviridae 03305

Super Taxa

DNA and RNA Reverse Transcribing Viruses; Viruses; Microorganisms

Organism Name

human immunodeficiency virus-1: pathogen

Taxa Notes

DNA and RNA Reverse Transcribing Viruses, Microorganisms, Viruses

L98 ANSWER 28 OF 47 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN

AN 2002:215999 BIOSIS

DN PREV200200215999

TI Infectious and whole **inactivated** simian immunodeficiency viruses interact similarly with primate **dendritic cells** (DCs): Differential intracellular fate of virions in mature and immature DCs.

AU Frank, I.; Piatak, M., Jr.; Stoessel, H.; Romani, N.; Bonnyay, D.; Lifson, J. D.; Pope, M. [Reprint author]

CS Population Council, Center for Biomedical Research, 1230 York Ave., New York, NY, 10021, USA
mpope@popcbr.rockefeller.edu

SO Journal of Virology, (March, 2002) Vol. 76, No. 6, pp. 2936-2951. print. CODEN: JOVIAM. ISSN: 0022-538X.

DT Article

LA English

ED Entered STN: 27 Mar 2002
Last Updated on STN: 27 Mar 2002

AB As potential targets for **human immunodeficiency virus** type 1 and simian immunodeficiency virus (HIV-1 and SIV), **dendritic cells** (DCs) likely play a significant role in the onset and spread of infection as well as in the induction of antiviral immunity. Using the SIV-macaque system to study the very early events in DC-virus interactions, we compared chemically **inactivated** SIV having conformationally and functionally intact envelope glycoproteins (2,2'-dithiodipyridine (AT-2) SIV) to infectious and heat-treated SIV. Both human and macaque DCs interact similarly with SIV without detectable effects on DC viability, phenotype, or endocytic function. As assessed by measuring cell-associated viral RNA, considerable amounts of virus are captured by the DCs and this is reduced when the virus is heat treated or derived from a strain that expresses low levels of envelope glycoprotein. Immunostaining for SIV proteins and electron microscopy indicated that few intact virus particles are retained at the periphery of the endocytically active, immature DCs. This contrasts with a perinuclear localization of numerous virions in large vesicular compartments deeper within mature DCs (in which macropinocytosis is down-regulated). Both immature and mature DCs are capable of clathrin-coated pit-mediated uptake of SIV, supporting the notion that the receptor-mediated uptake of virus can occur readily in mature DCs. While large numbers of whole viruses were preferentially found in mature DCs, both immature and mature DCs contained similar amounts of viral RNA, suggesting that different uptake/virus entry mechanisms are active in immature and mature DCs. These findings have significant implications for cell-to-cell transmission of HIV-1 and SIV and support the use of AT-2 SIV, an authentic but noninfectious form of virus, as a useful tool for studies of processing and presentation of AT-2 SIV antigens by DCs.

CC Cytology - Animal 02506
Biochemistry studies - Nucleic acids, purines and pyrimidines 10062
Virology - Animal host viruses 33506
Immunology - General and methods 34502
Medical and clinical microbiology - Virology 36006

IT Major Concepts

Immune System (Chemical Coordination and Homeostasis); Infection

IT Parts, Structures, & Systems of Organisms
 dendritic cell: immune system, differential
 intracellular fate, immature, mature, viability

IT Chemicals & Biochemicals
 2,2'-dithiodipyridine[AT-2]; RNA

IT Methods & Equipment
 electron microscopy: analytical method; immunostaining: analytical
 method

ORGN Classifier
 Cercopithecidae 86205
 Super Taxa
 Primates; Mammalia; Vertebrata; Chordata; Animalia
 Organism Name
 Macaca mulatta [rhesus macaque]: female, male
 Taxa Notes
 Animals, Chordates, Mammals, Nonhuman Mammals, Nonhuman Vertebrates,
 Nonhuman Primates, Primates, Vertebrates

ORGN Classifier
 Retroviridae 03305
 Super Taxa
 DNA and RNA Reverse Transcribing Viruses; Viruses; Microorganisms
 Organism Name
 human immunodeficiency virus type 1 [
 HIV-1]: cell-to-cell transmission
 simian immunodeficiency virus [SIV]: **inactivated**, infectious,
 whole
 Taxa Notes
 DNA and RNA Reverse Transcribing Viruses, Microorganisms, Viruses

L98 ANSWER 29 OF 47 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN
 AN 2002:625479 BIOSIS
 DN PREV200200625479
 TI Protection from SIV disease by vaccination with autologous dendritic cells
 pulsed with AT-2-inactivated whole virus.
 AU Zhu, Yong-de [Reprint author]; Koo, Kevin; Sutton, William F. [Reprint
 author]; Kuller, LaRene; Hu, Shiu-Lok; Benveniste, Raoul; Thomas, Elaine
 K.; Lifson, Jeffrey D.; Haigwood, Nancy L.
 CS Seattle Biomedical Research Institute, Seattle, WA, USA
 SO Journal of Medical Primatology, (August, 2002) Vol. 31, No. 4-5, pp.
 305-306. print.
 Meeting Info.: 19th Annual Symposium on Nonhuman Primate Models for AIDS.
 Monterey, CA, USA. September 08-11, 2002.
 CODEN: JMPMAO. ISSN: 0047-2565.

DT Conference; (Meeting)
 Conference; Abstract; (Meeting Abstract)

LA English

ED Entered STN: 12 Dec 2002
 Last Updated on STN: 12 Dec 2002

CC General biology - Symposia, transactions and proceedings 00520
 Cytology - Animal 02506
 Biochemistry studies - Proteins, peptides and amino acids 10064
 Pathology - Therapy 12512
 Blood - Blood and lymph studies 15002
 Blood - Blood cell studies 15004
 Endocrine - General 17002
 Pharmacology - General 22002
 Virology - Animal host viruses 33506
 Immunology - General and methods 34502
 Immunology - Immunopathology, tissue immunology 34508
 Medical and clinical microbiology - Virology 36006

IT Major Concepts
 Immune System (Chemical Coordination and Homeostasis); Infection;
 Pharmacology

IT Parts, Structures, & Systems of Organisms
 B cell: blood and lymphatics, immune system; CD4-positive cell: blood and lymphatics, immune system; T cell: blood and lymphatics, immune system; monocyte-derived dendritic cell: blood and lymphatics, immune system

IT Diseases
 SIV infection: immune system disease, viral disease, simian immunodeficiency virus infection
 Simian Acquired Immunodeficiency Syndrome (MeSH)

IT Chemicals & Biochemicals
 GM-CSF [granulocyte-macrophage colony stimulating factor]; IL-4 [interleukin-4]; SIV antigens; autologous dendritic cells pulsed with AT-2-inactivated whole virus vaccine

IT Methods & Equipment
 vaccination: immunologic method

IT Miscellaneous Descriptors
 immune response; viral titer; Meeting Abstract

ORGN Classifier
 Cercopithecidae 86205
 Super Taxa
 Primates; Mammalia; Vertebrata; Chordata; Animalia
 Organism Name
 Macaca fascicularis: host
 Taxa Notes
 Animals, Chordates, Mammals, Nonhuman Mammals, Nonhuman Vertebrates, Nonhuman Primates, Primates, Vertebrates

ORGN Classifier
 Retroviridae 03305
 Super Taxa
 DNA and RNA Reverse Transcribing Viruses; Viruses; Microorganisms
 Organism Name
 SIV [simian immunodeficiency virus]: mne, pathogen
 Taxa Notes
 DNA and RNA Reverse Transcribing Viruses, Microorganisms, Viruses

RN 83869-56-1 (GM-CSF)
 83869-56-1 (granulocyte-macrophage colony stimulating factor)

L98 ANSWER 35 OF 47 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN
 AN 2001:542898 BIOSIS
 DN PREV200100542898
 TI In vitro **HIV** eradication by autologous CD8+ T cells expanded with **inactivated-virus-pulsed dendritic cells**

AU Lu, Wei [Reprint author]; Andrieu, Jean-Marie [Reprint author]
 CS Laboratoire d'Oncologie et Virologie Moléculaire, Faculté Necker, Centre Biomedical des Saints-Peres, Paris, France
 SO Journal of Human Virology, (May-June, 2001) Vol. 4, No. 3, pp. 131. print.
 Meeting Info.: 2001 International Meeting of the Institute of Human Virology. Baltimore, Maryland, USA. September 09-13, 2001. Institute of Human Virology.
 ISSN: 1090-9508.

DT Conference; (Meeting)
 Conference; Abstract; (Meeting Abstract)

LA English

ED Entered STN: 21 Nov 2001
 Last Updated on STN: 25 Feb 2002

CC General biology - Symposia, transactions and proceedings 00520
 Cytology - Animal 02506
 Cytology - Human 02508
 Pathology - General 12502
 Pathology - Therapy 12512
 Blood - Blood and lymph studies 15002
 Blood - Blood cell studies 15004

Virology - Animal host viruses 33506
 Immunology - General and methods 34502
 Immunology - Immunopathology, tissue immunology 34508
 Medical and clinical microbiology - Virology 36006

IT Major Concepts
 Human Medicine (Medical Sciences); Immune System (Chemical Coordination and Homeostasis); Infection; Methods and Techniques

IT Parts, Structures, & Systems of Organisms
 CD8 T cells: blood and lymphatics, immune system; PBMC: blood and lymphatics, immune system, peripheral blood mononuclear cell;
 dendritic cells: immune system, **inactivated**
 virus-pulsed; lymphoid tissue: blood and lymphatics; monocyte: blood and lymphatics, immune system

IT Diseases
 HIV infection: immune system disease, viral disease,
 human immunodeficiency virus infection
 HIV Infections (MeSH)

IT Chemicals & Biochemicals
 adrithiol-2

IT Methods & Equipment
 highly active antiretroviral therapy: therapeutic method

IT Miscellaneous Descriptors
 Meeting Abstract

ORGN Classifier
 Hominidae 86215
 Super Taxa
 Primates; Mammalia; Vertebrata; Chordata; Animalia
 Organism Name
 human: host
 Taxa Notes
 Animals, Chordates, Humans, Mammals, Primates, Vertebrates

ORGN Classifier
 Retroviridae 03305
 Super Taxa
 DNA and RNA Reverse Transcribing Viruses; Viruses; Microorganisms
 Organism Name
 HIV [human immunodeficiency virus
]: pathogen
 Taxa Notes
 DNA and RNA Reverse Transcribing Viruses, Microorganisms, Viruses

L98 ANSWER 36 OF 47 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN
 AN 2000:330891 BIOSIS
 DN PREV200000330891
 TI Enhanced binding of antibodies to neutralization epitopes following thermal and chemical **inactivation** of **human immunodeficiency virus** type 1.
 AU Grovit-Ferbas, K.; Hsu, J. F.; Ferbas, J.; Gudeman, V.; Chen, I. S. Y.
 [Reprint author]
 CS UCLA School of Medicine, 11-934 Factor Building, Los Angeles, CA, 90095, USA
 SO Journal of Virology, (July, 2000) Vol. 74, No. 13, pp. 5802-5809. print. CODEN: JOVIAM. ISSN: 0022-538X.
 DT Article
 LA English
 ED Entered STN: 2 Aug 2000
 Last Updated on STN: 7 Jan 2002
 AB **Inactivation** of viral particles is the basis for several vaccines currently in use. Initial attempts to use simian immunodeficiency virus to model a killed **human immunodeficiency virus** type 1 (HIV-1) vaccine were unsuccessful, and limited subsequent effort has been directed toward a systematic study of the requirements for a protective killed HIV

-1 vaccine. Recent insights into HIV-1 virion and glycoprotein structure and neutralization epitopes led us to revisit whether **inactivated HIV-1** particles could serve as the basis for an HIV-1 vaccine. Our results indicate that relatively simple processes involving thermal and chemical **inactivation** can **inactivate HIV-1** by at least 7 logs. For some HIV-1 strains, significant amounts of envelope glycoproteins are retained in high-molecular-weight fractions. Importantly, we demonstrate retention of each of three conformation-dependent neutralization epitopes. Moreover, reactivity of monoclonal antibodies directed toward these epitopes is increased following treatment, suggesting greater exposure of the epitopes. In contrast, treatment of free envelope under the same conditions leads only to decreased antibody recognition. These **inactivated** virions can also be presented by human **dendritic cells** to direct a cell-mediated immune response in vitro. These data indicate that a systematic study of **HIV-1 inactivation**, gp120 retention, and epitope reactivity with conformation-specific neutralizing antibodies can provide important insights for the development of an effective killed **HIV-1** vaccine.

CC Biochemistry studies - Proteins, peptides and amino acids 10064
 Biochemistry studies - Carbohydrates 10068
 Pharmacology - Immunological processes and allergy 22018
 Virology - Animal host viruses 33506
 Immunology - General and methods 34502

IT Major Concepts
 Pharmacology

IT Parts, Structures, & Systems of Organisms
dendritic cells: immune system

IT Chemicals & Biochemicals
 anti-neutralization epitopes monoclonal antibodies: binding;
 conformation-dependent neutralization epitopes; envelope glycoproteins;
 gp120: glycoprotein; high-molecular-weight fractions; **human immunodeficiency virus-1** vaccine:
 immunostimulant-drug; killed **human immunodeficiency virus** type 1 vaccine: immunostimulant-drug; neutralization epitopes

IT Miscellaneous Descriptors
 cell-mediated immune response; vaccine development

ORGN Classifier
Retroviridae 03305
 Super Taxa
 DNA and RNA Reverse Transcribing Viruses; Viruses; Microorganisms
 Organism Name
human immunodeficiency virus-1; chemical
inactivation, thermal **inactivation**
 Taxa Notes
 DNA and RNA Reverse Transcribing Viruses, Microorganisms, Viruses

=> d his

(FILE 'HOME' ENTERED AT 10:36:24 ON 03 FEB 2004)
 SET COST OFF

FILE 'HCAPLUS' ENTERED AT 10:36:34 ON 03 FEB 2004
 E DENDRITIC CELL/CT

E E3+ALL
 L1 529 S E8,E9
 L2 7326 S E7
 L3 45 S E13
 L4 7836 S L1-L3
 L5 11672 S DENDRITIC CELL

L6 11672 S L4,L5
 E HIV/CT
 E E3+ALL
 L7 9029 S E2
 E E6+ALL
 L8 12470 S E7,E8,E9,E10
 L9 19620 S E6
 E E5+ALL
 L10 16484 S E6
 L11 36735 S E5+NT
 L12 635 S L6 AND L7-L11
 L13 639 S L6 AND HIV
 L14 613 S L6 AND HUMAN IMMUNODEFICIEN? VIRUS
 L15 798 S L12-L14
 L16 32 S L15 AND INACTIV?
 L17 58 S L15 AND PULS?
 L18 7 S L16 AND L17
 E CD8/CT
 E E10+ALL
 L19 8218 S E20
 L20 84 S L15 AND L19
 L21 150 S L15 AND CD8
 L22 150 S L20,L21
 L23 20 S L22 AND L16,L17
 L24 3 S L18 AND L23
 L25 4 S L18 NOT L24
 L26 7 S L24,L25
 L27 17 S L23 NOT L26
 SEL DN AN 6 14 15 16 17
 L28 5 S E1-E15 AND L27
 L29 12 S L26,L28
 L30 35 S L17 NOT L23-L29
 SEL DN AN 25
 L31 1 S E16-E18
 L32 13 S L29,L31
 L33 1 S US20040009194/PN OR US2002-390625#/AP,PRN
 E ANDRIEU J/AU
 L34 95 S E3,E6,E7,E12,E13,E17
 E LU L/AU
 L35 437 S E3-E26
 E LU LOUIS/AU
 L36 5 S E3,E4
 L37 4 S L15 AND L34-L36
 L38 4 S L34 AND L35-L36
 L39 19 S L32,L33,L37,L38
 L40 15 S L39 AND ?ACTIV?
 L41 4 S L39 NOT L40

FILE 'HCAPLUS' ENTERED AT 10:59:40 ON 03 FEB 2004

FILE 'MEDLINE' ENTERED AT 11:04:22 ON 03 FEB 2004

 E DENDRITIC CELL/CT
 E E5+ALL
 L42 9140 S E10
 L43 14071 S DENDRITIC CELL
 L44 14071 S L42,L43
 L45 878 S L44 AND HIV
 E HIV/CT
 E E3+ALL
 L46 564 S L44 AND E14+NT
 E HIV/CT
 L47 162 S L44 AND (E53+NT OR E28+NT)
 L48 0 S L44 AND E114+NT

L49 38 S L44 AND E89+NT
 L50 63 S L44 AND E149+NT
 L51 10 S L44 AND E191+NT
 L52 0 S L44 AND E216+NT
 E E231+ALL
 L53 492 S L44 AND E16+NT
 E E46+ALL
 L54 3 S L44 AND E14+NT
 E HIV INFECTION/CT
 L55 13 S L44 AND E76+NT
 L56 11 S L44 AND (E102+NT OR E104+NT)
 L57 9 S L44 AND E166+NT
 L58 31 S L44 AND E173+NT
 L59 0 S L44 AND E206+NT
 L60 0 S L44 AND E230+NT
 E E257+ALL
 E E23+ALL
 L61 138 S L44 AND E20+NT
 L62 37 S L44 AND (E41+NT OR E42+NT OR E43+NT OR E44+NT OR E45+NT)
 L63 320 S L44 AND HUMAN IMMUNODEFICIEN? VIRUS
 L64 927 S L45-L63
 L65 29 S L64 AND INACTIV?
 L66 64 S L64 AND PULS?
 E CD8/CT
 E E12+ALL
 L67 134 S L64 AND E17+NT
 L68 29 S L67 AND L65,L66
 L69 7 S L65 AND L66
 L70 3 S L68 AND L69
 E IMMUNITY/CT
 E E12+ALL
 L71 173712 S E4+NT
 L72 203 S L71 AND L64
 L73 9 S L72 AND L65
 L74 24 S L72 AND L66
 L75 63 S L72 AND L67
 L76 12 S L75 AND L73-L74
 L77 12 S L70,L76
 L78 26 S L69,L73,L65 NOT L77
 SEL DN AN 1 2
 L79 2 S L78 AND E1-E6
 L80 14 S L77,L79
 L81 14 S L80 AND L42-L80
 E ANDRIEU J/AU
 L82 208 S E3,E5,E8,E9
 E LU L/AU
 L83 833 S E3-E26
 E LU LOUIS/AU
 L84 1 S E3
 L85 63 S L82,L83,L84 AND L44
 L86 3 S L85 AND L64
 L87 15 S L81,L86

FILE 'MEDLINE' ENTERED AT 11:19:27 ON 03 FEB 2004

FILE 'BIOSIS' ENTERED AT 11:19:41 ON 03 FEB 2004

L88 16406 S L43
 L89 959 S L88 AND (HIV OR HUMAN() (IMMUNODEFICIEN? OR IMMUN# DEFICIEN?))
 L90 1048 S L88 AND RETROVIRIDAE+NT/BC
 L91 1048 S L88 AND RETROVIRIDAE+NT/ORGAN
 L92 215 S L90,L91 NOT L89
 L93 12 S L92 AND INACTIV?
 SEL DN AN 5 12

L94 2 S E1-E4
L95 833 S L89 AND L90,L91
L96 28 S L95 AND INACTIV?
L97 30 S L94,L96

FILE 'HCAPLUS, MEDLINE, BIOSIS' ENTERED AT 11:24:53 ON 03 FEB 2004
L98 47 DUP REM L40 L87 L97 (13 DUPLICATES REMOVED)

=>

THOMSON
MICROMEDEX
PDR® Electronic Library™

PDR SUITE | INTERACTIONS | STEDMAN'S DICTIONARY

DICTIONARY SEARCH | BROWSE

Return to
Thomson Subscription Page

Log Out

[Dictionary Search](#) > Definition



STEDMAN'S The Best Words In Medicine.™

Stedman's Dictionary

Define:

Search

Stedman's Medical Dictionary 27th Edition

autologous (aw-to'l'o-gus)

1. Occurring naturally and normally in a certain type of tissue or in a specific structure of the body. 2. In transplantation, referring to a graft in which the donor and recipient areas are in the same individual, or to blood that the donor has previously donated and then receives back, usually during surgery. 3. Sometimes used to denote a neoplasm derived from cells that occur normally at that sight, e.g., a squamous cell carcinoma in the upper esophagus. SYN: autogenous (1) . [auto- + G. logos, 1 relation]

Copyright© 2000 Lippincott Williams & Wilkins. All rights reserved.

[Home](#) || [Contact Us](#) || [Warranty and Disclaimer](#) || [About Us](#) || [Help](#) || [Log Out](#)
Copyright© 2002-2004 Thomson PDR. All rights reserved.

BEST AVAILABLE COPY



BEST AVAILABLE COPY